

Asymptotics for minimisers of convex processes

Nils Lid Hjort and David Pollard

University of Oslo and Yale University

ABSTRACT. By means of two simple convexity arguments we are able to develop a general method for proving consistency and asymptotic normality of estimators that are defined by minimisation of convex criterion functions. This method is then applied to a fair range of different statistical estimation problems, including Cox regression, logistic and Poisson regression, least absolute deviation regression outside model conditions, and pseudo-likelihood estimation for Markov chains.

Our paper has two aims. The first is to exposit the method itself, which in many cases, under reasonable regularity conditions, leads to new proofs that are simpler than the traditional proofs. Our second aim is to exploit the method to its limits for logistic regression and Cox regression, where we seek asymptotic results under as weak regularity conditions as possible. For Cox regression in particular we are able to weaken previously published regularity conditions substantially.

KEY WORDS: *argmin lemma approximation, convexity, Cox regression, LAD regression, log-concavity, logistic regression, minimal conditions, partial likelihood, pseudo-likelihood*

1. Introduction. This paper develops a simple method for proving consistency and asymptotic normality for estimators defined by minimisation of a convex criterion function. Versions of the method have been used or partially used by several authors, for various specific occasions, including Jurečková (1977, 1991), Andersen and Gill (1982), Hjort (1986, 1988a), Haberman (1989), Pollard (1990, 1991), Bickel, Klassen, Ritov and Wellner (1992), Niemiro (1992), but the general principle has not been widely recognised.

Our aims in this paper are twofold. (i) The primary objective is to explain the basic method, and to illustrate its use in a fair range of statistical estimation problems. In section 2 we state and prove some general theorems about estimators that are defined via some form of convex minimisation, and in sections 3 and 4 illustrate their use by means of applications to sample quantiles, maximum likelihood estimation when the likelihood is log-concave, and least squares and least absolute deviation linear regression outside model conditions. Similarly sections 5 and 6 treat logistic and Cox regression, while still further applications are reported in section 7, including Poisson regression. The proofs are relatively simple and instructive, at least when regularity conditions are kept reasonable. (ii) The second objective is to improve on previously published results, in the sense of pruning down the regularity conditions of theorems for two important models, namely logistic regression in section 5 and Cox regression in sections 6 and 7A. The two aims are mildly conflicting, editorially speaking. We soften the conflict in sections 5 and 6 by writing down first a simple version of a theorem with a simple proof, and then a harder version with a harder proof. In this way we hope that our article has some pedagogic merits while at the same time also offering something to the specialists.

Instead of treating minimisation as a search for a root of a derivative, we work directly with the argmin (a minimising value) of a random function and are able to approximate it with the argmin of a simpler random function. In this way we manage to avoid special arguments that are often used to prove consistency separately. Convexity essentially buys us both consistency and asymptotic normality with the same dollar, and sometimes with cheaper regularity conditions.

The two convexity lemmas that will be used are as follows.

LEMMA 1: FROM POINTWISE TO UNIFORM. Suppose $A_n(s)$ is a sequence of convex random functions defined on an open convex set \mathcal{S} in \mathbb{R}^p , which converges in probability to some $A(s)$, for each s . Then $\sup_{s \in K} |A_n(s) - A(s)|$ goes to zero in probability, for each compact subset K of \mathcal{S} .

PROOF: This is proved in Andersen and Gill (1982, appendix), crediting T. Brown, via ‘diagonal subsequencing’ and an appeal to a corresponding non-stochastic result (see Rockafellar, 1970, Theorem 10.8). For a direct proof, see Pollard (1991, section 6). \square

A convex function is continuous and attains its minimum on compact sets, but it can be flat at its bottom and have several minima. For simplicity we speak about ‘the argmin’ when referring to any of the possible minimisers. The argmin can be selected in a measurable way, as explained in Niemiro (1992, p. 1531), for example.

LEMMA 2: NEARNESS OF ARGMINS. Suppose $A_n(s)$ is convex as in Lemma 1 and is approximated by $B_n(s)$. Let α_n be the argmin of A_n , and assume that B_n has a unique argmin β_n . Then there is a probabilistic bound on how far α_n can be from β_n : for each $\delta > 0$,

$$\Pr\{|\alpha_n - \beta_n| \geq \delta\} \leq \Pr\{\Delta_n(\delta) \geq \frac{1}{2}h_n(\delta)\}, \quad (1.1)$$

where

$$\Delta_n(\delta) = \sup_{|s - \beta_n| \leq \delta} |A_n(s) - B_n(s)| \quad \text{and} \quad h_n(\delta) = \inf_{|s - \beta_n| = \delta} B_n(s) - B_n(\beta_n). \quad (1.2)$$

PROOF: The lemma as stated has nothing to do with convergence or indeed with the ‘ n ’ subscript at all, of course, but is stated in a form useful for later purposes. To prove it, let s be an arbitrary point outside the ball around β_n with radius δ , say $s = \beta_n + lu$ for a unit vector u , where $l > \delta$. Convexity of A_n implies

$$(1 - \delta/l) A_n(\beta_n) + (\delta/l) A_n(s) \geq A_n(\beta_n + \delta u).$$

Writing for convenience $A_n(s) = B_n(s) + r_n(s)$, we deduce

$$\begin{aligned} (\delta/l) \{A_n(s) - A_n(\beta_n)\} &\geq A_n(\beta_n + \delta u) - A_n(\beta_n) \\ &= B_n(\beta_n + \delta u) + r_n(\beta_n + \delta u) - B_n(\beta_n) - r_n(\beta_n) \\ &\geq h_n(\delta) - 2\Delta_n(\delta). \end{aligned}$$

If $\Delta_n(\delta) < \frac{1}{2}h_n(\delta)$, then $A_n(s) > A_n(\beta_n)$ for all s outside the δ -ball, which means that the minimiser α_n must be inside. This proves (1.1).

It is worth pointing out that any norm on \mathbb{R}^p can be used here, and that no assumptions need to be placed on the B_n function beside the existence of the minimiser β_n . \square

The two lemmas will deliver more than mere consistency when applied to suitably rescaled and recentred versions of convex processes.

We record a couple of useful implications of Lemma 2. If $A_n - B_n$ goes to zero uniformly on bounded sets in probability and β_n is stochastically bounded, then $\Delta_n(\delta) \rightarrow_p 0$ by a simple argument. It follows that $\alpha_n - \beta_n \rightarrow_p 0$ provided only that $1/h_n(\delta)$ is stochastically bounded for each fixed δ . This last requirement says that B_n shouldn’t flatten out around its minimum as n increases.

BASIC COROLLARY. Suppose $A_n(s)$ is convex and can be represented as $\frac{1}{2}s'Vs + U_n's + C_n + r_n(s)$, where V is symmetric and positive definite, U_n is stochastically bounded, C_n is arbitrary, and $r_n(s)$ goes to zero in probability for each s . Then α_n , the argmin of A_n , is only $o_p(1)$ away from $\beta_n = -V^{-1}U_n$, the argmin of $\frac{1}{2}s'Vs + U_n's + C_n$. If also $U_n \rightarrow_d U$ then $\alpha_n \rightarrow_d -V^{-1}U$.

PROOF: The function $A_n(s) - U'_n s - C_n$ is convex and goes to $\frac{1}{2}s'Vs$ in probability for each s . By the first lemma the convergence is uniform on bounded sets. Let $\Delta_n(\delta)$ be the supremum of $|r_n(s)|$ over $\{|s - \beta_n| \leq \delta\}$. Then, by Lemma 2,

$$\alpha_n = -V^{-1}U_n + \varepsilon_n, \quad \text{where } \Pr\{|\varepsilon_n| \geq \delta\} \leq \Pr\{\Delta_n(\delta) \geq \frac{1}{2}k\delta^2\} \rightarrow 0. \quad (1.3)$$

Here k is the smallest eigenvalue of V , and $\Delta_n(\delta) \rightarrow_p 0$, by the arguments used above. \square

A useful slight extension of this is when $A_n(s) = \frac{1}{2}s'V_n s + U'_n s + C_n + r_n(s)$ is convex, with a nonnegative definite symmetric V_n matrix that converges in probability to a positive definite V . Writing $V_n = V + \eta_n$ the remainder η_n can be absorbed into $r_n(s)$ and the result above holds.

2. General results for convex minimisation estimators. This section presents three basic theorems about the asymptotic behaviour of estimators that are defined by minimisation of some convex criterion function. The first is for the independent identically distributed (i.i.d.) case. The second is stated for independent observations with different distributions, and is suitable for proving consistency and asymptotic normality in regression models, for example, under model conditions. The third theorem also applies to regression model estimators, but is suited to give asymptotic results also outside model conditions. Applications and illustrations are provided in sections 3, 4 and 5.

2A. A theorem for the i.i.d. case. Let Y_1, Y_2, \dots be i.i.d. from some distribution F . A certain p -dimensional parameter $\theta_0 = \theta(F)$ is of interest. Assume that one of the ways of characterising this parameter is to say that it minimises $Eg(Y, t) = \int g(y, t) dF(y)$, where the $g(y, t)$ function is convex in t . Examples include quantiles, the mean, M-estimation and maximum likelihood estimation parameters and so on; see sections 3 and 4. In the expectation expression above, and later on, Y denotes a generic observation from the true underlying F .

Some weak expansion of $g(y, t)$ around the value θ_0 of t is needed, but we avoid explicitly requiring pointwise derivatives to exist. With this in mind, write

$$g(y, \theta_0 + t) - g(y, \theta_0) = D(y)'t + R(y, t) \quad (2.1)$$

for a $D(y)$ with mean zero under F . If $ER(Y, t)^2$ is of order $o(|t|^2)$ as $t \rightarrow 0$, as we will usually require, then $D(y)$ is nothing but the derivative in quadratic mean of the function $g(y, \theta_0 + t)$ at $t = 0$.

THEOREM 2.1. *Suppose that $g(y, t)$ is convex in t as above, and that (2.1) holds with*

$$E\{g(Y, \theta_0 + t) - g(Y, \theta_0)\} = ER(Y, t) = \frac{1}{2}t'Jt + o(|t|^2) \text{ as } t \rightarrow 0 \quad (2.2)$$

for a positive definite matrix J . Suppose also that $\text{Var } R(Y, t) = o(|t|^2)$, and that $D(Y)$ has a finite covariance matrix $K = \int D(y)D(y)' dF(y)$. Then the estimator $\hat{\theta}_n$ which minimises $\sum_{i \leq n} g(Y_i, t)$ is \sqrt{n} -consistent for θ_0 , and

$$\sqrt{n}(\hat{\theta}_n - \theta_0) = -J^{-1}n^{-1/2} \sum_{i \leq n} D(Y_i) + o_p(1). \quad (2.3)$$

In particular $\sqrt{n}(\hat{\theta}_n - \theta_0) \rightarrow_d -J^{-1}\mathcal{N}_p\{0, K\} = \mathcal{N}_p\{0, J^{-1}KJ^{-1}\}$.

PROOF: Consider the convex function $A_n(s) = \sum_{i \leq n} \{g(Y_i, \theta_0 + s/\sqrt{n}) - g(Y_i, \theta_0)\}$. It is minimised by $\sqrt{n}(\hat{\theta}_n - \theta_0)$. Note first that $nER(Y, s/\sqrt{n}) = \frac{1}{2}s'Js + r_{n,0}(s)$ where $r_{n,0}(s) = no(|s|^2/n) \rightarrow 0$ for fixed s . Accordingly, using (2.1),

$$\begin{aligned} A_n(s) &= \sum_{i \leq n} \{D(Y_i)'s/\sqrt{n} + R(Y_i, s/\sqrt{n}) - ER(Y_i, s/\sqrt{n})\} + nER(Y, s/\sqrt{n}) \\ &= U_n's + \frac{1}{2}s'Js + r_{n,0}(s) + r_n(s), \end{aligned}$$

in which

$$U_n = n^{-1/2} \sum_{i \leq n} D(Y_i) \quad \text{and} \quad r_n(s) = \sum_{i \leq n} \{R(Y_i, s/\sqrt{n}) - ER(Y_i, s/\sqrt{n})\}.$$

Now $r_n(s)$ tends to zero in probability for each s , since its mean is zero and its variance is $\sum_{i \leq n} \text{Var} R(Y_i, s/\sqrt{n}) = no(1/n)$. This, together with the Basic Corollary of section 1, proves (2.3) and the limit distribution result, since U_n goes to a $\mathcal{N}_p\{0, K\}$ by the central limit theorem. Note that both consistency and asymptotic normality followed from the same approximation argument. \square

Note that $\text{Var} R(Y, t) = ER(Y, t)^2 + O(t^4)$, so we might as well work with second moments rather than variances. Notice also that the differentiability assumption (2.2) is applied to the process obtained by averaging out over the distribution F , a smoothing that can eliminate troublesome pointwise behaviour of $R(y, t)$. Huber (1967) recognised this advantage of smoothing before differentiating.

2B. A theorem for independent observations with different distributions. Assume that the true density of Y_i is of the form $f_i(y_i) = f_i(y_i, \theta_0, \eta_i)$, where θ_0 is a certain p -dimensional parameter of interest. Suppose that an estimator $\hat{\theta}_n$ for θ_0 is proposed which minimises $\sum_{i \leq n} g_i(Y_i, \theta)$, where the $g_i(y_i, \theta)$ functions are convex in θ . A simple example is linear regression, where $Y_i = \theta_0'x_i + \varepsilon_i$ and $\hat{\theta}_n$ minimises $\sum_{i \leq n} (Y_i - \theta'x_i)^2$.

Suppose that $g_i(y_i, \theta_0 + t) - g_i(y_i, \theta_0) = D_i(y_i)'t + R_i(y_i, t)$, where $ED_i(Y_i) = 0$. With the previous development in mind, write

$$ER_i(Y_i, t) = \frac{1}{2}t'A_it + v_{i,0}(t) \quad \text{and} \quad \text{Var} R_i(Y_i, t) = v_i(t), \quad (2.4)$$

and let B_i be the variance matrix for $D_i(Y_i)$. The sums $J_n = \sum_{i \leq n} A_i$ and $K_n = \sum_{i \leq n} B_i$ are featured below. The first useful result, properly generalising Theorem 2.1, is the following, which is proved by copying the arguments of 2A mutatis mutandis.

THEOREM 2.2. *Assume that $\sum_{i \leq n} v_{i,0}(s/\sqrt{n}) \rightarrow 0$ and $\sum_{i \leq n} v_i(s/\sqrt{n}) \rightarrow 0$ for each s , and that J_n/n and K_n/n converge to J and K , where J is positive definite. Then $\sqrt{n}(\hat{\theta}_n - \theta_0)$ is only $o_p(1)$ away from $-J^{-1}n^{-1/2} \sum_{i \leq n} D_i(Y_i)$. If in particular the Lindeberg requirements are fulfilled for the $D_i(Y_i)$ sequence, then $\sqrt{n}(\hat{\theta}_n - \theta_0) \rightarrow_d \mathcal{N}_p\{0, J^{-1}KJ^{-1}\}$.*

Another result which sometimes is stronger is as follows. Assume that $\sum_{i \leq n} v_{i,0}(J_n^{-1/2}s) \rightarrow 0$ and $\sum_{i \leq n} v_i(J_n^{-1/2}s) \rightarrow 0$ for each s , and that $J_n^{-1}K_n$ is bounded. Then

$$J_n^{1/2}(\hat{\theta}_n - \theta_0) = -J_n^{-1/2}K_n^{1/2}U_n + o_p(1), \quad (2.5)$$

where $U_n = K_n^{-1/2} \sum_{i \leq n} D_i(Y_i)$. If in particular there are matrices J and K such that $J_n^{-1}K_n$ goes to $J^{-1}K$, and the Lindeberg conditions are fulfilled, securing $U_n \rightarrow_d \mathcal{N}_p\{0, I_p\}$, then $J_n^{1/2}(\hat{\theta}_n -$

$\theta_0\} \rightarrow_d \mathcal{N}_p\{0, J^{-1/2}KJ^{-1/2}\}$. This result is proved by studying the convex function $\sum_{i \leq n} \{g_i(Y_i, \theta_0 + J_n^{-1/2}s) - g_i(Y_i, \theta_0)\}$. In some situations of interest $J_n = K_n$, further simplifying (2.5). See section 5 for an illustration of this.

2C. A theorem for regression type estimators outside model conditions. The results of 2B are sometimes not sufficient. Theorem 2.3 below will work for asymptotic behaviour of regression methods outside model conditions, as made clear in section 3D, for example.

Assume that some covariate vector $x_i = (x_{i,1}, \dots, x_{i,p})'$ is associated with observation Y_i . For simplicity we formulate a result in terms of densities, rather than general distribution functions. Suppose that the true density for Y_i given x_i is $f(y_i|x_i)$ but that some regression model postulates $f(y_i, \beta|x_i)$, for a suitable p -dimensional parameter vector β . We consider an estimator $\hat{\beta}_n$ defined to minimise $\sum_{i \leq n} g_i(Y_i, \beta|x_i)$, where $g_i(y_i, \beta|x_i)$ is convex in β for each (y_i, x_i) . In the following we shall assume that the empirical distribution of x_1, \dots, x_n , whether actually random or under the experimenter's control, converges to a well-defined distribution H in x -space. This conceptual limit is to be thought of as the 'covariate distribution'. Assume that $n^{-1} \sum_{i \leq n} g_i(Y_i, \beta|X_i)$ converges in probability to a function with a unique minimiser β_0 .

Under these circumstances it is not generally possible to get a representation like the one that led to (2.4), because of heterogeneity as well as potential modelling bias, as the applications in section 3D and section 5C will illustrate. It becomes necessary to include a x_i -dependent bias term. Suppose that it is possible to write

$$g_i(y_i, \beta_0 + t|x_i) - g_i(y_i, \beta_0|x_i) = \{\delta(x_i) + D_i(y_i|x_i)\}'t + R_i(y_i, t|x_i), \quad (2.6)$$

where $ED_i(Y_i|x_i) = 0$ and $\text{VAR} D_i(Y_i|x_i) = B_i(x_i)$. Write furthermore

$$ER_i(Y_i, t|x_i) = \frac{1}{2}t'A_i(x_i)t + v_{i,0}(t|x_i) \quad \text{and} \quad \text{Var} R_i(Y_i, t|x_i) = v_i(t|x_i). \quad (2.7)$$

This time three matrix sums are needed, $J_n = \sum_{i \leq n} A_i(x_i)$, $K_n = \sum_{i \leq n} B_i(x_i)$, and $L_n = \sum_{i \leq n} \delta(x_i)\delta(x_i)'$.

THEOREM 2.3. *Assume that the x_1, x_2, \dots sequence is such that*

$$\sum_{i \leq n} v_{i,0}(s/\sqrt{n}|x_i) \rightarrow 0 \quad \text{and} \quad \sum_{i \leq n} v_i(s/\sqrt{n}|x_i) \rightarrow_p 0 \quad \text{for each } s, \quad (2.8)$$

that the J_n/n sequence is bounded away from zero, and that the K_n/n and L_n/n sequences are bounded. Then

$$\sqrt{n}(\hat{\beta}_n - \beta_0) = -(J_n/n)^{-1} \left\{ n^{-1/2} \sum_{i \leq n} \delta(x_i) + n^{-1/2} \sum_{i \leq n} D_i(Y_i|x_i) \right\} + \varepsilon_n, \quad (2.9)$$

where $\varepsilon_n = \varepsilon_n(x_1, \dots, x_n) \rightarrow_p 0$.

The proof is quite similar to previous proofs in this section, taking as its starting point the convex function $\sum_{i \leq n} \{g_i(Y_i, \beta_0 + s/\sqrt{n}|x_i) - g_i(Y_i, \beta_0|x_i)\}$. We omit the details.

The (2.9) representation has two statistically interesting implications. (i) In the conditional framework with a given x_i sequence, suppose that $J_n/n \rightarrow J$ and $K_n/n \rightarrow K$ and that the Lindeberg condition holds for $\sum_{i \leq n} n^{-1/2} D_i(Y_i|x_i)$. Then

$$\sqrt{n}(\hat{\beta}_n - \beta_0)|x_1, \dots, x_n = \mathcal{N}_p \left\{ -(J_n/n)^{-1} n^{-1/2} \sum_{i \leq n} \delta(x_i), J^{-1}KJ^{-1} \right\} + \varepsilon_n', \quad (2.10)$$

where $\varepsilon'_n \rightarrow_p 0$. So $\widehat{\beta}_n$ is approximately normal with variance matrix $J^{-1}KJ^{-1}/n$, but actually biased with a bias depending on x_1, \dots, x_n . The bias is typically zero under exact regression model conditions, see 3D below. (ii) Secondly, if the x_i 's can be treated as being independent and coming from their own 'design distribution' $H(dx)$ in x -space, then $\delta(x_i)$ has mean zero and variance matrix L , say. In this unconditional framework

$$\sqrt{n}(\widehat{\beta}_n - \beta_0) = J^{-1}\mathcal{N}_p\{0, K + L\} + o_p(1) \rightarrow_d \mathcal{N}_p\{0, J^{-1}(K + L)J^{-1}\}. \quad (2.11)$$

3. Applications and illustrations.

3A. The median. Let Y_1, Y_2, \dots be i.i.d. from a density f , let μ be the population median, and let M_n be the sample median from the first n observations. We shall prove the well known fact that M_n is consistent for μ and that

$$\sqrt{n}(M_n - \mu) \rightarrow_d \mathcal{N}\{0, 1/4f(\mu)^2\}, \quad (3.1)$$

provided only that f is positive and continuous at μ .

This fits into the framework of 2A with the convex function $g(y, t) = |y - t|$. The (2.1) expansion reads

$$|y - (\mu + t)| - |y - \mu| = D(y)t + R(y, t),$$

where $D(y) = -I\{y > \mu\} + I\{y \leq \mu\}$, and

$$R(y, t) = \begin{cases} 2(t - (y - \mu))I\{\mu \leq y \leq \mu + t\} & \text{if } t > 0, \\ 2((y - \mu) - t)I\{\mu + t \leq y \leq \mu\} & \text{if } t < 0, \end{cases}$$

while $R(y, 0) = 0$, which makes it easy to verify

$$ER(Y, t) = f(\mu)t^2 + o(t^2) \quad \text{and} \quad ER(Y, t)^2 = \frac{4}{3}f(\mu)|t|^3 + o(|t|^3).$$

Actually we only need a distribution function with a positive derivative at μ . Of course we don't get the explicit $|t|^3$ bound then. Notice that $D(Y)$ and $R(Y, t)$ are bounded functions even if $|Y - t|$ itself can have infinite expected value, since we work with the difference $|Y - (\mu + t)| - |Y - \mu|$. Since the variance of $D(Y)$ is equal to 1, assertion (3.1) follows from Theorem 2.1. See 4A below for an extension of this result.

3B. Simultaneous asymptotic normality of order statistics. Let f be positive and continuous in its support region, and consider the function

$$g_p(y, t) = p\{(y - t)_+ - y_+\} + (1 - p)\{(t - y)_+ - (-y)_+\}.$$

It is convex in t and its expected value is minimal for $t = F^{-1}(p) = \mu_p$, the p -th quantile of the underlying distribution, and

$$E\{g_p(Y, t) - g_p(Y, \mu_p)\} = \frac{1}{2}f(\mu_p)(t - \mu_p)^2 + o((t - \mu_p)^2)$$

can be shown. The (2.1) expansion works with

$$D(y) = (1 - p)I\{y \leq \mu_p\} - pI\{y > \mu_p\} = I\{y \leq \mu_p\} - p$$

and $ER(Y, t)^2 = O(|t|^3)$ can be checked. Let $Q_{n,p}$ be the minimiser of $\sum_{i \leq n} g_p(Y_i, t)$, which is sometimes non-unique, but which in any case is at most $O_p(n^{-1})$ away from the $[np]$ 'th order statistic $Y_{([np])}$. The general theorem of 2A implies

$$Z_n(p) = \sqrt{n}(Q_{n,p} - \mu_p) = -f(\mu_p)^{-1} \sqrt{n}\{F_n(\mu_p) - p\} + \varepsilon_n(p), \quad (3.2)$$

where F_n is the empirical distribution function and $\varepsilon_n(p) \rightarrow 0$ in probability for each p . This links the quantile process Z_n to the empirical process, and proves finite-dimensional convergence in distribution of the quantile process to a Gaussian process $Z(\cdot)$ with mean zero and covariance structure

$$\text{cov}\{Z(p_1), Z(p_2)\} = \frac{p_1(1-p_2)}{f(\mu_{p_1})f(\mu_{p_2})} \quad \text{for } p_1 \leq p_2. \quad (3.3)$$

The traditional proofs of this finite-dimensional convergence result are rather messier than the above. There is in reality also process convergence here, of course, which is linked to the fact that $\sup_{\delta \leq s \leq 1-\delta} |\varepsilon_n(p)|$ goes to zero in probability for each δ . Proving this is not within easy reach of our method, however. See also the comment ending 3D below.

3C. Estimation in L_α mode. Let more generally $M_{n,\alpha}$ minimise $\sum_{i \leq n} |Y_i - t|^\alpha$, where $\alpha \geq 1$, and let ξ_α be the population parameter that minimises $E|Y - t|^\alpha$. For $\alpha = \frac{3}{2}$ we would expect an estimator with properties somehow between those for the median and the mean, for example. We can prove

$$\sqrt{n}(M_{n,\alpha} - \xi_\alpha) \rightarrow_d \mathcal{N}\{0, \tau^2\} \quad \text{where } \tau^2 = \frac{E|Y - \xi_\alpha|^{2(\alpha-1)}}{\{(\alpha-1)E|Y - \xi_\alpha|^{\alpha-2}\}^2}, \quad (3.4)$$

assuming $E|Y|^{2(\alpha-1)}$ to be finite. The proof proceeds by mimicking that for the simpler case $\alpha = 1$. One needs to use

$$D(y) = -\alpha(y - \xi_\alpha)^{\alpha-1} I\{y > \xi_\alpha\} + \alpha(\xi_\alpha - y)^{\alpha-1} I\{y < \xi_\alpha\},$$

and it is somewhat more cumbersome but feasible to bound $ER(Y, t)^2$. And finally needed is the analytical fact that $E\{|Y - (\xi_\alpha + t)|^\alpha - |Y - \xi_\alpha|^\alpha\} = \frac{1}{2} K_f t^2 + o(t^2)$, in which $K_f = \alpha(\alpha-1)E|Y - \xi_\alpha|^{\alpha-2}$.

It is interesting to note here that $(\alpha-1)E|Y - \xi_\alpha|^{\alpha-2}$ tends to $2f(F^{-1}(\frac{1}{2}))$ as α tends to 1, explaining the connection from the moment-type expression for the variance τ^2 of (3.4) to the rather different-looking expression for the median case.

It is also worth pointing out that the (3.4) result can be reached via influence functions and function space methods as well. The influence function can be found to be

$$I(F, y) = \begin{cases} \alpha K_F^{-1} |y - \xi_\alpha(F)|^{\alpha-1} & \text{if } y > \xi_\alpha(F), \\ -\alpha K_F^{-1} |y - \xi_\alpha(F)|^{\alpha-1} & \text{if } y < \xi_\alpha(F), \end{cases}$$

after which the usual argument is that since $\sqrt{n}(M_{n,\alpha} - \xi_\alpha) = n^{-1/2} \sum_{i \leq n} I(F, Y_i) + \varepsilon_n$, for suitable remainder term ε_n , one must have limiting normality with $\tau^2 = \int I(F, y)^2 dF(y)$, agreeing with (3.4). But proving that ε_n here goes to zero in probability is not trivial, since the ξ_α functional is rather non-smooth. The argument can be saved via establishing Lipschitz differentiability, as in Example 1 of Huber (1967). Our method manages to avoid these somewhat sophisticated arguments.

3D. Agnostic least squares and least absolute deviation regression. Statistical regression is about estimating the unknown centre value of Y for given x , i.e. the curve or surface centre($Y|x$),

based on $p + 1$ -tuplets (x_i, Y_i) , where ‘centre’ could be the mean or the median. Ordinary linear regression uses a linear approximation $\beta'x = \sum_{j=1}^p \beta_j x_j$ for this centre function, which is often a very reasonable method even if the true underlying centre function is somewhat non-linear. The least squares regression estimator is $\hat{\beta}'_n x$ where $\hat{\beta}_n$ minimises $\sum_{i \leq n} (Y_i - \beta'x_i)^2$, and the least absolute deviation estimator is $\tilde{\beta}'_n x$ where $\tilde{\beta}_n$ minimises $\sum_{i \leq n} |Y_i - \beta'x_i|$.

Statistical properties of these estimators are usually investigated only under the admittedly unlikely assumption that the true surface *is* linear and that the variances are constant over the full region, i.e.

$$Y_i = \beta'_0 x_i + \sigma \varepsilon_i \quad (3.5)$$

where the ε_i 's are i.i.d. standardised residuals centred around zero. An in some sense more honest approach would be to merely postulate that

$$Y_i = m(x_i) + \sigma(x_i)\varepsilon_i, \quad (3.6)$$

for some smooth functions $m(x)$ and $\sigma(x)$, and view the regression surface estimator as an attempt to produce a good linear approximation to the evasive $m(x)$. Our plan now is to derive properties under robust and agnostic (3.6) conditions using Theorem 2.3 of 2C, while assuming that the empirical distribution of x_i 's converges to an appropriate ‘covariate distribution’ H . Under ideal (3.5) conditions they specialise to results obtainable using the simpler Theorem 2.2 of 2B.

Consider least squares regression first, assuming the ε_i 's to have mean zero and variance one. This fits into the 2C framework with $g_i(Y_i, \beta|x_i) = \frac{1}{2}(Y_i - \beta'x_i)^2$. The method aims at getting the best linear approximation $\beta'_0 x$ to $m(x)$, in the sense of minimising the limit of $n^{-1} \sum_{i \leq n} (m(x_i) - \beta'x_i)^2$. In fact this means $\beta_0 = (EXX')^{-1}EXY$. We find

$$\begin{aligned} g_i(Y_i, \beta_0 + t|x_i) - g_i(Y_i, \beta_0|x_i) &= -(Y_i - \beta'_0 x_i)x'_i t + \frac{1}{2}(t'x_i)^2 \\ &= -(\delta(x_i) + D_i(Y_i|x_i))'t + \frac{1}{2}t'x_i x'_i t, \end{aligned}$$

in which

$$\delta(x_i) = (m(x_i) - \beta'_0 x_i)x_i \quad \text{and} \quad D_i(Y_i|x_i) = (Y_i - m(x_i))x_i.$$

In the notation of (2.7) one has $A_i(x_i) = x_i x'_i$ and both remainder terms are simply equal to zero. Consider

$$J_n = \sum_{i \leq n} x_i x'_i, \quad K_n = \sum_{i \leq n} \sigma(x_i)^2 x_i x'_i, \quad L_n = \sum_{i \leq n} \{m(x_i) - \beta'_0 x_i\}^2 x_i x'_i. \quad (3.7)$$

Two results can be given, corresponding to (2.10) and (2.11). First, suppose the x_i sequence is such that $J_n/n \rightarrow$ a positive definite J , $K_n/n \rightarrow K$, that the L_n/n sequence is bounded, and that $\max_{i \leq n} \sigma(x_i)^2 |x_i|^2 / \sum_{i \leq n} \sigma(x_i)^2 |x_i|^2 \rightarrow 0$. Then $\sqrt{n}(\hat{\beta}_n - \beta_0)$ is asymptotically normal with mean $J^{-1}n^{-1/2} \sum_{i \leq n} (m(x_i) - \beta'_0 x_i)x_i$ and variance matrix $J^{-1}KJ^{-1}$. Secondly, under the unconditional viewpoint where the x_i 's are seen as i.i.d. with finite variance matrix $L = E(m(X) - \beta'_0 X)^2 X X'$ for $\delta(x_i)$, then

$$\sqrt{n}(\hat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_p\{0, J^{-1}(K + L)J^{-1}\}. \quad (3.8)$$

Note that $K + L$ can be estimated consistently with $n^{-1} \sum_{i \leq n} (Y_i - \hat{\beta}'_n x_i)^2 x_i x'_i$.

These results can also be derived more or less directly, i.e. without the convex machinery of section 2, see Exercise 45 in Hjort (1988b). In the least absolute deviation case to be reported on next a direct approach is much more difficult, however, but it can be efficiently handled using the methods of section 2.

For the LAD regression case, take the ε_i 's of (3.6) to have distribution F with median zero and variance one. We will assume that F has a density f which further possesses a continuous derivative f' . In this case $g_i(Y_i, \beta|x_i) = |Y_i - \beta'x_i|$, and the method aims at getting the best approximation β'_0x to $m(x)$ in the sense of minimising the long term value of $n^{-1} \sum_{i \leq n} \mathbb{E}|m(x_i) - \beta'x_i + \sigma(x_i)\varepsilon_i|$. We skip the various details that have to be worked through to reach a result here. They resemble those above and arguments used in 3A. To give the result, let $D_i(Y_i|x_i) = 1$ if $Y_i \leq \beta'_0x_i$ and -1 if $Y_i > \beta'_0x_i$, with conditional mean $h(x_i) = 2 \Pr\{m(x_i) + \sigma(x_i)\varepsilon_i \leq \beta'_0x_i\} - 1$, and consider the three matrices

$$J_n = \sum_{i \leq n} 2f_i(\beta'_0x_i - m(x_i))x_ix_i', \quad K_n = \sum_{i \leq n} \{1 - h(x_i)\}^2 x_ix_i', \quad L_n = \sum_{i \leq n} h(x_i)^2 x_ix_i',$$

where $f_i(z) = f(z/\sigma(x_i))/\sigma(x_i)$ is the density of the scaled residual $\sigma(x_i)\varepsilon_i$. In particular $K_n + L_n = \sum_{i \leq n} x_ix_i'$. As for the least squares case these efforts lead to a representation

$$\sqrt{n}(\tilde{\beta}_n - \beta_0) = -(J_n/n)^{-1} \left[n^{-1/2} \sum_{i \leq n} h(x_i)x_i + n^{-1/2} \sum_{i \leq n} \{D_i(Y_i|x_i) - h(x_i)\}x_i \right] + \varepsilon_n. \quad (3.9)$$

This has one implication for given x_i -sequences and another implication for the 'overall variability'. Under some mild assumptions $J_n/n \rightarrow J$ and $(K_n + L_n)/n \rightarrow K + L$, and $\sqrt{n}(\tilde{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_p\{0, J^{-1}(K + L)J^{-1}\}$. The $K + L$ matrix is estimated consistently using $\sum_{i \leq n} x_ix_i'/n$ whereas a more complicated consistent estimate, involving smoothing and density estimation, can be constructed for J .

The special case of $\text{med}(Y|x) = m(x) = \beta'_0x$ has $J_n = \sum_{i \leq n} 2f_i(0)x_ix_i'/\sigma(x_i)$, and the perfect but perhaps unrealistic case of both a linear median and a constant variance has $J_n^{-1}(K_n + L_n)J_n^{-1} = \{4f(0)^2\}^{-1}(\sum_{i \leq n} x_ix_i')^{-1}\sigma^2$. This is the case considered in Pollard (1990).

Our method can also be applied to the quantile regression situation, where one aims to estimate $m(x_0) + \sigma(x_0)F^{-1}(p)$, for example, to construct a prediction interval for a future Y at a given covariate value x_0 . This time one minimises $\sum_{i \leq n} g_p(Y_i, \beta'x_i)$ with the g_p function of 3B. This gives a suitable generalisation of results reached by Bassett and Koenker (1982).

4. Maximum likelihood type estimation.

4A. *Log-concave densities.* Suppose Y_1, Y_2, \dots are i.i.d. from some continuous density f , and that a parametric model of the form $f(y, \theta) = f(y, \theta_1, \dots, \theta_p)$ is employed, where the parameter space is some open and convex region. We stipulate that $\log f(y, \theta)$ be concave in θ in this region and shall be able to reprove familiar results on maximum likelihood (ML) and Bayes estimation, using the convexity based results of section 2, but with milder smoothness assumptions than those traditionally employed.

Note that the log-likelihood $\sum_{i \leq n} \log f(Y_i, \theta)$ when divided by n tends to $\mathbb{E} \log f(Y, \theta) = \int f(y) \log f(y, \theta) dy$, for each θ . Assume that this function has a unique global maximum at θ_0 , which is the 'agnostic parameter value' that gives best approximation according to the Kullback-Leibler distance $\int f(y) \log\{f(y)/f(y, \theta)\} dy$ from truth to approximating density. From section 2A the following result is quite immediate.

THEOREM 4.1. *Suppose $\log f(y, \theta_0 + t) - \log f(y, \theta_0) = D(y)'t + R(y, t)$ is concave in t , for a $D(\cdot)$ function with mean zero and finite covariance matrix K under f , and that the remainder term satisfies*

$$\mathbb{E}\{\log f(Y, \theta_0 + t) - \log f(Y, \theta_0)\} = \mathbb{E}R(Y, t) = -\frac{1}{2}t'Jt + o(|t|^2) \quad (4.1)$$

as well as $\text{Var } R(Y_i, t) = o(|t|^2)$, where J is symmetric and positive definite. Then the maximum likelihood estimator $\hat{\theta}_n$ is \sqrt{n} -consistent for θ_0 and

$$\sqrt{n}(\hat{\theta}_n - \theta_0) = J^{-1}n^{-1/2} \sum_{i \leq n} D(Y_i) + o_p(1) \rightarrow_d J^{-1} \mathcal{N}\{0, K\} = \mathcal{N}_p\{0, J^{-1}KJ^{-1}\}.$$

In ordinary smooth cases one can Taylor expand and use $D(y) = \partial \log f(y, \theta_0) / \partial \theta$ and find a remainder $R(y, t)$ with mean $-\frac{1}{2}t'Jt + O(|t|^3)$ and squared mean of order $O(|t|^4)$, involving

$$J = -E_f \frac{\partial^2 \log f(Y_i, \theta_0)}{\partial \theta \partial \theta} \quad \text{and} \quad K = \text{VAR}_f \frac{\partial \log f(Y_i, \theta_0)}{\partial \theta}. \quad (4.2)$$

Notice that when the model happens to be perfect, as in textbooks for optimistic statisticians, then $K = J$, and we get the more familiar $\mathcal{N}_p\{0, J^{-1}\}$ result.

EXAMPLE. In addition to the median M_n in the situation of 3A, look at the mean absolute deviation statistic $\hat{\tau}_n = n^{-1} \sum_{i \leq n} |Y_i - M_n|$. We will show simultaneous convergence of $\sqrt{n}(M_n - \mu, \hat{\tau}_n - \tau)$, where $\tau = E|Y_i - \mu|$, and for this assume finite variance of the Y_i 's.

This can be accomplished by considering the parametric model $f(y, \mu, \tau) = (2\tau)^{-1} \exp\{-|y - \mu|/\tau\}$ for data. This model may be quite inadequate to describe the behaviour of the data sequence, but the ML estimates are nevertheless M_n and $\hat{\tau}_n$ as above. The traditional theorems on ML behaviour require more smoothness than is present here, and indeed often require that the true f belongs to the model, but Theorem 4.1 can be used. This is because $\log f(y, \mu, \tau)$ is concave in $(\mu, 1/\tau)$. Verifying conditions involves details similar to those in 3A, and we omit them here. The result is

$$\begin{pmatrix} \sqrt{n}(M_n - \mu) \\ \sqrt{n}(\hat{\tau}_n - \tau) \end{pmatrix} \rightarrow_d \mathcal{N}_2 \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1/\{4f(\mu)^2\}, & \text{cov} \\ \text{cov}, & \text{Var } Y_i - \tau^2 \end{pmatrix} \right\}.$$

where the covariance is $E\{Y_i \leq \mu\} |Y_i - \mu| - \frac{1}{2}\tau$. Note that there is asymptotic independence if f is symmetric around μ .

4B. *Bayes and maximum likelihood estimators are asymptotically equivalent.* It is well known that Bayes and ML estimation are asymptotically equivalent procedures in regular situations. In other words, if θ_n^* is the Bayes estimator under some prior $\pi(\theta)$, then $\sqrt{n}(\theta_n^* - \theta_0)$ has the same limit distribution as $\sqrt{n}(\hat{\theta}_n - \theta_0)$. The standard proofs of this fact involve many technicalities, and furthermore are typically restricted to calculations under the assumption that the underlying $f(y, \theta_0)$ model is exactly correct, see e.g. Lehmann (1983, chapter 6.7). Below follows a reasonably quick proof of this fact, and it is reassuring that the result is valid also outside model circumstances.

Let $\pi(\theta)$ be a prior density, assumed continuous at θ_0 and satisfying the growth constraint

$$\pi(\theta) \leq C_1 \exp(C_2|\theta|) \quad \text{for all } \theta,$$

where C_1 and C_2 are positive constants. The posterior density is proportional to $L_n(\theta)\pi(\theta)$, where $L_n(\theta) = \prod_{i \leq n} f(Y_i, \theta)$ is the likelihood. The Bayes estimator θ_n^* (under quadratic loss) is the posterior mean. Note that improper priors are accepted too.

We shall make use of the following dominated convergence fact, which is a special case of Lemma A3 in the appendix. Suppose $\{G_n(s, \omega)\}$ is a sequence of random functions (assumed jointly measurable) such that $G_n(s, \omega) \rightarrow G(s)$ in probability, for each s . Suppose $H(s)$ is an integrable function for which the set $\{\omega: |G_n(s, \omega)| \leq H(s) \text{ for all } s\}$ has probability tending to one. Then $\int G_n(s, \omega) ds \rightarrow \int G(s) ds$ in probability. (Apply Lemma A3 with X_n equal to G_n restricted to the set where $G_n \leq H$.)

THEOREM 4.2. *Under the conditions of Theorem 4.1, the MLE estimator $\widehat{\theta}_n$ and the posterior mean θ_n^* are asymptotically equivalent, in the sense that $\sqrt{n}(\theta_n^* - \widehat{\theta}_n) \rightarrow_p 0$.*

PROOF: Define the random convex function $A_n(s)$ by

$$\exp(-A_n(s)) = L_n(\widehat{\theta}_n + s/\sqrt{n})/L_n(\widehat{\theta}_n).$$

By definition of the ML estimator, A_n achieves its minimum value of zero at $s = 0$. By the change of variables $\theta = \widehat{\theta}_n + s/\sqrt{n}$ we find

$$\theta_n^* = \frac{\int \theta L_n(\theta) \pi(\theta) d\theta}{\int L_n(\theta) \pi(\theta) d\theta} = \widehat{\theta}_n + \frac{1}{\sqrt{n}} \frac{\int s \exp(-A_n(s)) \pi(\widehat{\theta}_n + s/\sqrt{n}) \exp(-C_2|\widehat{\theta}_n|) ds}{\int \exp(-A_n(s)) \pi(\widehat{\theta}_n + s/\sqrt{n}) \exp(-C_2|\widehat{\theta}_n|) ds}.$$

The random function A_n converges in probability uniformly on compact sets to $\frac{1}{2}s'Js$. Define $\gamma_n = \inf_{|t|=1} A_n(t)$. It converges in probability to $\gamma_0 = \inf_{|t|=1} \frac{1}{2}t'Jt > 0$. Argue as in Lemma 2 to show that $A_n(s) \geq \gamma_n|s|$ for $|s| > 1$. The domination condition needed for the fact noted above holds in both numerator and denominator with

$$H(s) = \begin{cases} 2C_1 & \text{if } |s| \leq 1, \\ C_1|s| \exp(-\frac{1}{2}\gamma_0|s|) & \text{if } |s| > 1. \end{cases}$$

The ratio of integrals converges in probability to

$$\frac{\int s \exp(-\frac{1}{2}s'Js) \pi(\theta_0) \exp(-C_2|\theta_0|) ds}{\int \exp(-\frac{1}{2}s'Js) \pi(\theta_0) \exp(-C_2|\theta_0|) ds} = 0.$$

The result follows. \square

5. Logistic regression. Suppose that $p + 1$ -tuplets (x_i, Y_i) are observed, where $x_i = (x_{i,1}, \dots, x_{i,p})'$ is a covariate vector ‘explaining’ the binomial outcome Y_i . The logistic regression model postulates that the Y_i ’s are independent with

$$\Pr\{Y_i = 1|x_i\} = q(x_i, \beta) = \frac{\exp(\beta'x_i)}{1 + \exp(\beta'x_i)} \quad \text{for some } \beta = \beta_0, \quad (5.1)$$

and the ML estimator $\widehat{\beta}_n = (\widehat{\beta}_{n,1}, \dots, \widehat{\beta}_{n,p})'$ maximises the log-likelihood function

$$\sum_{i \leq n} [Y_i \log q(x_i, \beta) + (1 - Y_i) \log\{1 - q(x_i, \beta)\}] = \sum_{i \leq n} [Y_i \beta'x_i - \log\{1 + \exp(\beta'x_i)\}].$$

Of course the asymptotic normality of this estimator is well known and widely used, but precise sufficient conditions are not easy to find in the literature.

We will soon arrive at such, employing results of 2B, which are applicable since the summands above are concave in β . As a preparatory exercise we mark down the following little expansion, which holds for all u and $u + h$, in terms of $\pi(u) = \exp(u)/\{1 + \exp(u)\}$:

$$\log \frac{1 + \exp(u + h)}{1 + \exp(u)} = \pi(u)h + \frac{1}{2}\pi(u)\{1 - \pi(u)\}h^2 + \frac{1}{6}\pi(u)\{1 - \pi(u)\}\gamma(u, h)h^3, \quad (5.2)$$

where $|\gamma(u, h)| \leq \exp(|h|)$. This is proved from the exact third order Taylor expansion expression, with third term equal to $\frac{1}{6}\pi(u')\{1 - \pi(u')\}\{1 - 2\pi(u')\}h^3$, for appropriate u' between u and $u + h$.

Some analysis reveals that $\pi(u')\{1 - \pi(u')\} \leq \exp(|h|) \pi(u)\{1 - \pi(u)\}$, regardless of u and h . This is in fact quite similar to what results from using Lemma A2 in the appendix, but the bound on the remainder obtained here suits the problem better.

5A. Under model conditions. In the spirit of our two aims, laid out in the Introduction, we will first give a simpler result with a ‘pedagogical proof’, and then sharpen the tools to reach a second result with minimal regularity conditions. Under model conditions (5.1), write for convenience $q_i = q(x_i, \beta_0)$, and let $J_n = \sum_{i \leq n} q_i(1 - q_i)x_i x_i'$ be the information matrix.

THEOREM 5.1. *Assume that $\mu_n = \max_{i \leq n} |x_i|/\sqrt{n} \rightarrow 0$ and that $J_n/n \rightarrow J$. Then, under model conditions (5.1), $\sqrt{n}(\hat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_p\{0, J^{-1}\}$.*

PROOF: We will use Theorem 2.2 with $g_i(y_i, \beta) = \log f_i(y_i, \beta) = y_i \beta' x_i - \log\{1 + \exp(\beta' x_i)\}$. The expansion noted above yields

$$\begin{aligned} \log \frac{f_i(y_i, \beta_0 + t)}{f_i(y_i, \beta_0)} &= y_i t' x_i - [\log\{1 + \exp(\beta_0' x_i + t' x_i)\} - \log\{1 + \exp(\beta_0' x_i)\}] \\ &= (y_i - q_i)x_i' t - \frac{1}{2}q_i(1 - q_i)(t' x_i)^2 - \frac{1}{6}q_i(1 - q_i)\gamma_i(t)(t' x_i)^3 \\ &= D_i(y_i)' t - R_i(y_i, t). \end{aligned}$$

Here $D_i(y_i) = (y_i - q_i)x_i$ and $R_i(y_i, t) = \frac{1}{2}t' q_i(1 - q_i)x_i x_i' t + v_{i,0}(t)$, where $|\gamma_i(t)| \leq \exp(|t' x_i|)$ in the expression for $v_{i,0}(t)$. Note that $J_n = K_n$, in the notation of Theorem 2.2, and that $R_i(Y_i, t)$ has zero variance, so what we have to prove is (i) that $\sum_{i \leq n} v_{i,0}(s/\sqrt{n}) \rightarrow 0$, (ii) that the Lindeberg conditions are satisfied for $\sum_{i \leq n} n^{-1/2}(Y_i - q_i)x_i$. But

$$\begin{aligned} \left| \sum_{i \leq n} v_{i,0}(s/\sqrt{n}) \right| &\leq \sum_{i \leq n} \frac{1}{6}q_i(1 - q_i) \exp(|s' x_i/\sqrt{n}|) |s' x_i/\sqrt{n}|^3 \\ &\leq \sum_{i \leq n} \frac{1}{6}q_i(1 - q_i) \exp(|s|\mu_n) (s' x_i x_i' s/n) |s|\mu_n \\ &= \frac{1}{6}|s|\mu_n \exp(|s|\mu_n) s'(J_n/n)s, \end{aligned}$$

which goes to zero. And the Lindeberg condition is that for each s and δ

$$\sum_{i \leq n} E n^{-1}(Y_i - q_i)^2 (s' x_i)^2 I\{|(Y_i - q_i)s' x_i/\sqrt{n}| \geq \delta\} \rightarrow 0,$$

and this sum is bounded by $s'(J_n/n)s I\{|s|\mu_n| \geq \delta\}$. This ends the proof. \square

If the x_i 's are i.i.d. from some covariate distribution H , then $\mu_n \rightarrow 0$ a.s. exactly when the components of x_i have finite second moment. This also secures convergence of J_n/n to $J = \int q(x, \beta_0)\{1 - q(x, \beta_0)\} x x' H(dx)$.

Our second and sharper theorem is proved next, by squeezing more out of the bound of the $v_{i,0}(t)$ remainder and more out of the Lindeberg condition.

THEOREM 5.2. *Assume that the $\lambda_n = \max_{i \leq n} |J_n^{-1/2} x_i|$ sequence is bounded, and that*

$$N_n(\delta) = \sum_{i \leq n} q_i(1 - q_i) x_i' J_n^{-1} x_i I\{|J_n^{-1/2} x_i| \geq \delta\} \rightarrow 0 \quad \text{for each positive } \delta. \quad (5.3)$$

Then, under model conditions (5.1), $J_n^{1/2}(\hat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_p\{0, I_p\}$.

PROOF: We consider the random convex function $\sum_{i \leq n} \{\log f_i(Y_i, \beta_0 + J_n^{-1/2}s) - \log f_i(Y_i, \beta_0)\}$, which upon using the expansion again can be rearranged as $U'_n s - \frac{1}{2}s's - r_n(s)$, where $U_n = J_n^{-1/2} \sum_{i \leq n} (Y_i - q_i)x_i$ and $r_n(s) = \sum_{i \leq n} \frac{1}{6}q_i(1 - q_i)\gamma_i(s'J_n^{-1/2}x_i)(s'J_n^{-1/2}x_i)^3$. We are to prove (i) that $r_n(s) \rightarrow 0$, and (ii) that $U_n \rightarrow_d \mathcal{N}_p\{0, I_p\}$.

At this stage we call on appendix A1 where it is shown that (5.3) is a sufficient and actually also a necessary condition (ii) to hold. And $|r_n(s)|$ is bounded by $\sum_{i \leq n} \frac{1}{6}q_i(1 - q_i) \exp(|s'J_n^{-1/2}x_i|)|s'J_n^{-1/2}x_i|^3$. We split this sum into $|J_n^{-1/2}x_i| < \delta$ summands and $|J_n^{-1/2}x_i| \geq \delta$ summands. The first sum is bounded by $\frac{1}{6}|s|^3\delta \exp(|s|\delta)$, and the second is bounded by $\frac{1}{6}|s|^3\lambda_n \exp(|s|\lambda_n) N_n(\delta)$. Letting $n \rightarrow \infty$ and $\delta \rightarrow 0$ afterwards shows that indeed $r_n(s) \rightarrow 0$. \square

It is worth noting that the $N_n(\delta) \rightarrow 0$ condition in the theorem serves two purposes: forcing an analytic remainder term towards zero, and securing uniform negligibility of individual terms in the large-sample distribution of $J_n^{-1/2} \sum_{i \leq n} D_i(Y_i)$, i.e. a normal limit. Note also that $\lambda_n \rightarrow 0$ suffices for the conclusion to hold, since $N_n(\delta) \leq p\lambda_n/\delta$.

5B. Outside model conditions. Let us next depart from the strict model assumption (5.1), which in most cases merely is intended to provide a reasonable approximation to some more complicated reality, and stipulate only that $\Pr\{Y = 1|x\} = q(x)$ for some true, underlying $q(x)$ function. Fitting the logistic regression equation makes sense still, and turns out to aim at achieving the best approximation $q(x, \beta)$ to the true $q(x)$, in a sense made precise as follows. Let

$$\Delta_x[q(x), q(x, \beta)] = q(x) \log \frac{q(x)}{q(x, \beta)} + \{1 - q(x)\} \log \frac{1 - q(x)}{1 - q(x, \beta)}$$

be the Kullback–Leibler distance from true binomial $(1, q(x))$ to modelled binomial $(1, q(x, \beta))$, and let $\Delta[q(\cdot), q(\cdot, \beta)] = \int \Delta_x[q(x), q(x, \beta)] H(dx)$ be the weighted distance between the true probability curve to the modelled probability curve, in which H again is the ‘covariate distribution’ for x ’s, as discussed in 2C. The following can now be proved using methods of 2C: ML estimation is \sqrt{n} -consistent for the value β_0 that minimises the weighted Kullback–Leibler distance Δ , and $\sqrt{n}(\hat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_d\{0, J^{-1}KJ^{-1}\}$, provided the two matrices

$$J = \mathbb{E} X X' q(X, \beta_0) \{1 - q(X, \beta_0)\} = \int x x' q(x, \beta_0) \{1 - q(x, \beta_0)\} H(dx),$$

$$K = \mathbb{E} X X' \{Y - q(X, \beta_0)\}^2 = \int x x' [q(x) \{1 - q(x)\} + \{q(x) - q(x, \beta_0)\}^2] H(dx)$$

are finite. This result was also obtained in Hjort (1988a), where various implications for statistical inference also are discussed.

6. Cox regression. In this section new proofs are presented for the consistency and asymptotic normality of the usual estimators in Cox’s famous semiparametric regression model for survival analysis data. The parametric Cox regression model is somewhat simpler, and is treated in 7A below. The regularity requirements we need turn out in both cases to be weaker than those earlier presented in the literature.

The most complete results and proofs in the literature for the basic large-sample properties of the estimators in this model are perhaps those of Andersen and Gill (1982) and Hjort (1992). Andersen and Gill obtain results under the conditions of the model, and with regularity conditions quite weaker than earlier i.i.d. type assumptions, whereas Hjort explores the large-sample behaviour also outside the conditions of the model. For a history of the Cox model and the various approaches to reach asymptotics results, see Andersen, Borgan, Gill & Keiding (1992, chapter VII).

Our present intention is to provide yet another proof, which in several ways is simpler and requires less involvement with the martingale techniques than the one of Andersen and Gill. As in the previous section we choose to present two theorems, reflecting our two aims explained in section 1. The first holds when the covariates are bounded, in which case the proof is quite transparent, and extra regularity conditions can be kept quite minimal. The second version is more sophisticated in that it tolerates unbounded covariates and weakens regularity conditions further.

The usual Cox regression model for possibly censored lifetimes with covariate information is as follows: The individuals have independent lifetimes T_1^0, \dots, T_n^0 , and the i -th has hazard rate

$$\lambda_i(s) = \lambda(s) \exp(\beta' z_i(s)) = \lambda(s) \exp(\beta_1 z_{i,1}(s) + \dots + \beta_p z_{i,p}(s)), \quad (6.1)$$

depending on that person's covariate vector $z_i(s)$, and involving some unspecified basis hazard rate $\lambda(s)$. As indicated the covariates are allowed to depend on time s , and they can be random processes, as long as they are previsible; $z_i(s)$ should only depend on information available at time $s-$ (for a full discussion of previsibility, or predictability, see Andersen et al. (1992, p. 65–66)). There is a possibly interfering censoring time C_i leaving just $T_i = \min\{T_i^0, C_i\}$ and $\delta_i = I\{T_i^0 \leq C_i\}$ to the statistician. Consider the at risk indicator function $Y_i(s) = I\{T_i \geq s\}$, which is left continuous and hence previsible, and the counting process N_i with mass δ_i at T_i , i.e. $dN_i(s) = I\{T_i \in [s, s+ds], \delta_i = 1\}$. The log partial likelihood can then be written

$$G_n(\beta) = \sum_{i \leq n} \int_0^L \{\beta' z_i(s) - \log R_n(s, \beta)\} dN_i(s), \quad (6.2)$$

featuring the empirical relative risk function $R_n(s, \beta) = \sum_{i \leq n} Y_i(s) \exp(\beta' z_i(s))$; see for example Andersen et al. (1992, chapter VII). It is assumed that data are collected on the finite time interval $[0, L]$ only. The *Cox estimator* is the value $\hat{\beta}_n$ that maximises the partial likelihood.

Lemma A2 of the appendix allows us an expansion for $\log R_n(s, \beta_0 + x)$, using $w_i = Y_i(s) \exp(\beta_0' z_i(s))$ and $a_i = z_i(s)'x$. The result is

$$\log R_n(s, \beta_0 + x) - \log R_n(s, \beta_0) = \bar{z}_n(s)'x + \frac{1}{2}x'V_n(s)x + v_n(x, s), \quad (6.3)$$

where

$$\bar{z}_n(s) = \sum_{i \leq n} p_{n,i}(s) z_i(s) \quad \text{and} \quad V_n(s) = \sum_{i \leq n} p_{n,i}(s) (z_i(s) - \bar{z}_n(s))(z_i(s) - \bar{z}_n(s))', \quad (6.4)$$

and $p_{n,i}(s) = Y_i(s) \exp(\beta_0' z_i(s)) / R_n(s, \beta_0)$. A bound for the remainder term in (6.3) is $|v_n(x, s)| \leq \frac{4}{3} \max_{i \leq n} |(z_i(s) - \bar{z}_n(s))'x|^3$. Observe that $\bar{z}_n(s)$ and $V_n(s)$ can be interpreted as the mean value and the variance matrix for $z_i(s)$, where this covariate vector is randomly selected among those at risk at time s with probabilities proportional to the relative risks $\exp(\beta_0' z_i(s))$.

All this leaves us suitably prepared for a theorem.

THEOREM 6.1. *Assume that the hazard rate for the i 'th individual follows the Cox model (6.1) with a true parameter β_0 and a continuous positive basis hazard $\lambda(s)$, and that the covariate processes $z_i(s)$ are previsible and uniformly bounded. Assume that*

$$J_n(s) = n^{-1} \sum_{i \leq n} Y_i(s) \exp(\beta_0' z_i(s)) (z_i(s) - \bar{z}_n(s))(z_i(s) - \bar{z}_n(s))' \rightarrow_p J(s) \quad (6.5)$$

for almost all s in $[0, L]$ and that $J = \int_0^L J(s)\lambda(s) ds$ is positive definite. Then $\widehat{\beta}_n$ is \sqrt{n} -consistent for β_0 and $\sqrt{n}(\widehat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_p\{0, J^{-1}\}$.

PROOF: As a simple consequence of earlier efforts we have

$$\begin{aligned} G_n^*(x) &= G_n(\beta_0 + x/\sqrt{n}) - G_n(\beta_0) \\ &= \sum_{i \leq n} \int_0^L [n^{-1/2}(z_i(s) - \bar{z}_n(s))'x - \frac{1}{2}n^{-1}x'V_n(s)x - v_n(x/\sqrt{n}, s)] dN_i(s) \\ &= U_n'x - \frac{1}{2}x'J_n^*x - r_n(x), \end{aligned} \quad (6.6)$$

where we write

$$U_n = n^{-1/2} \sum_{i \leq n} \int_0^L (z_i(s) - \bar{z}_n(s)) dN_i(s) \quad \text{and} \quad J_n^* = n^{-1} \int_0^L V_n(s) d\bar{N}_n(s), \quad (6.7)$$

using $\bar{N}_n(\cdot) = \sum_{i \leq n} N_i(\cdot)$ to denote the aggregated counting process for the data. The designated remainder term is $r_n(x) = \int_0^L v_n(x/\sqrt{n}, s) d\bar{N}_n(s)$, which goes to zero, since it is bounded by $\int_0^L \frac{4}{3}(2K)^3|x|^3/n^{3/2} d\bar{N}_n(s)$, which is $O(n^{-1/2})$. The K here is the absolute bound on the covariates. That the (6.6) function is concave in x is clear from the convexity of $\log R_n(s, \beta)$ in β . By the basic method of section 1 it only remains to show (i) that $J_n^* \rightarrow_p J$ and (ii) that $U_n \rightarrow_d \mathcal{N}_p\{0, J\}$.

At this stage we need some of the easier bits of the martingale representation and convergence theory for counting processes, but manage to avoid needing some of the more sophisticated inequalities and technicalities that have invariably been present in earlier rigorous proofs, like in Andersen and Gill (1982). The counting process N_i has compensator process $A_i(t) = \int_0^t Y_i(s) \exp(\beta_0' z_i(s)) d\Lambda(s)$, writing $d\Lambda(s) = \lambda(s) ds$. This means that $M_i(t) = N_i(t) - A_i(t)$ is a zero mean martingale, with increments $dM_i(s) = dN_i(s) - Y_i(s) \exp(\beta_0' z_i(s)) d\Lambda(s)$. One can show that $M_i(t)^2 - A_i(t)$ as well as $M_i(t)M_j(t)$ are martingales too, when $i \neq j$, which in martingale theory parlance means that M_i has variance process $\langle M_i, M_i \rangle(t) = A_i(t)$ and that they are orthogonal, i.e. $\langle M_i, M_j \rangle = 0$. See Andersen et al. (1992, chapter II), for example. Inserting $dN_i(s) = dM_i(s) + dA_i(s)$ in (6.7) leads to

$$J_n^* = \int_0^L J_n(s) d\Lambda(s) + n^{-1} \sum_{i \leq n} \int_0^L V_n(s) dM_i(s) \quad (6.8)$$

and

$$U_n = n^{-1/2} \sum_{i \leq n} \int_0^L (z_i(s) - \bar{z}_n(s)) dM_i(s), \quad (6.9)$$

in that two other terms cancel.

We are now in a position to prove (i) and (ii). Note that the first term of (6.8) goes to J in probability by boundedness of the integrand and Lemma A3 in the appendix. The second term is $O_p(n^{-1/2})$, which can be seen using boundedness of covariates in conjunction with the result

$$\mathbb{E} \left\{ \int_0^L \sum_{i \leq n} H_i(s) dM_i(s) \right\}^2 = \mathbb{E} \sum_{i \leq n} \int_0^L H_i(s)^2 d\langle M_i, M_i \rangle(s),$$

valid for previsible random functions H_i . This proves (i). To prove convergence in distribution of U_n we essentially use the version of Rebolledo's martingale central limit theorem given in Andersen and Gill (1982, appendix I). Its variance process converges properly,

$$\langle U_n, U_n \rangle(L) = n^{-1} \sum_{i \leq n} \int_0^L (z_i(s) - \bar{z}_n(s))(z_i(s) - \bar{z}_n(s))' d\langle M_i, M_i \rangle(s) = \int_0^L J_n(s) d\Lambda(s) \rightarrow_p J,$$

and the necessary Lindeberg type condition is also satisfied:

$$n^{-1} \sum_{i \leq n} \int_0^L |z_i(s) - \bar{z}_n(s)|^2 I\{n^{-1/2}|z_i(s) - \bar{z}_n(s)| \geq \varepsilon\} Y_i(s) \exp(\beta'_0 z_i(s)) d\Lambda(s) \rightarrow_p 0 \quad (6.10)$$

since the indicator function ends up being zero for all large n . \square

Next we present a stronger theorem with weaker conditions imposed. The proof is basically the same as for the previous result, but more is squeezed out of bounds for remainder terms and out of conditions for the martingale convergence to hold.

THEOREM 6.2. *Assume that the hazard rate for the i 'th individual follows the Cox model (6.1) with a true parameter β_0 and a continuous positive basis hazard $\lambda(s)$. Assume that $J_n(s)$ goes to some $J(s)$ in probability for almost all s , as in (6.5), and that $\int_0^L J_n(s)\lambda(s) ds \rightarrow_p J = \int_0^L J(s)\lambda(s) ds$, a positive definite matrix. Suppose finally that*

$$\mu_n(s) = n^{-1/2} \max_{i \leq n} |z_i(s) - \bar{z}_n(s)| \rightarrow_p 0 \quad \text{for almost each } s \quad (6.11)$$

and that $\max_{s \leq L} \mu_n(s)$ is stochastically bounded. Then again $\sqrt{n}(\hat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_p\{0, J^{-1}\}$.

PROOF: (6.6) and (6.7) still hold, and we plan to demonstrate (i) $r_n(x) \rightarrow_p 0$, (ii) $J_n^* \rightarrow_p J$, and (iii) $U_n \rightarrow_d \mathcal{N}_p\{0, J\}$.

(i) is proved by using the tighter bound for $v_n(x, s)$ of (6.3) available by employing Lemma A2, namely $\frac{2}{3}g(\max_{i \leq n} |(z_i(s) - \bar{z}_n(s))'x|) x'V_n(s)x$, for $g(u) = u \exp(2u + 4u^2)$. This leads to

$$|r_n(x)| \leq \int_0^L \frac{2}{3}g(\mu_n(s)|x|) x'V_n(s)x d\bar{N}_n(s)/n.$$

Split this into two terms, using $d\bar{N}_n(s) = R_n(s, \beta_0) d\Lambda(s) + \sum_{i \leq n} dM_i(s)$. The first of the resulting terms goes to zero in probability by assumptions on $J_n(s)$ and dominated convergence (appendix A3), and the other term is of smaller stochastic order. Secondly (ii) follows as in the previous proof, since the second term of (6.8) vanishes in probability, by variations of the same arguments. Finally two ingredients are needed to secure (iii). The first is $\langle U_n, U_n \rangle(L) \rightarrow_p J$, which holds by assumptions as in the previous proof. The second is a more elaborate demonstration of the Lindeberg type condition (6.10), now accomplished by bounding it with

$$\int_0^L \text{Tr}(J_n(s)) I\{\mu_n(s) \geq \varepsilon\} d\Lambda(s),$$

which goes to zero in probability by dominated convergence (the integrand goes pointwise to zero in probability and is dominated by $\text{Tr}(J_n(s))$, see appendix A3 again).

And all this combined with the Basic Corollary triumphantly implies that the argmax of the (6.6) function, which is $\sqrt{n}(\hat{\beta}_n - \beta_0)$, is only $o_p(1)$ away from the argmax of $U'_n x - \frac{1}{2}x'Jx$, which is $J^{-1}U_n$. This proves consistency and asymptotic normality. \square

REMARKS. (i) Usually one would have $V_n(s) \rightarrow_p V(s)$ and $n^{-1}R_n(s, \beta_0) \rightarrow_p R(s, \beta_0)$, say, so that $J_n(s) = V_n(s)R_n(s, \beta_0)/n \rightarrow_p J(s) = V(s)R(s, \beta_0)$; in particular $J = \int_0^L V(s)R(s, \beta_0) d\Lambda(s)$ in such cases, and this is the expression typically encountered for the inverse covariance matrix. (ii) The Andersen and Gill regularity requirements include rather strong uniform convergence statements, in both time s and β near β_0 . In the development above this would mean requiring

$$\sup_{s \in [0, L]} \sup_{\beta \in U(\beta_0)} \left| n^{-1} \sum_{i \leq n} Y_i(s) z_i(s) z_i(s)' \exp(\beta' z_i(s)) - J(s, \beta) \right| \rightarrow_p 0,$$

for example, for a suitable neighbourhood $U(\beta_0)$ and a suitable limit function $J(s, \beta)$. This contrasts sharply with our condition (6.5), which is only about β_0 , and is pointwise in s . Andersen and Gill also include various other asymptotic stability conditions, about uniform continuity and differentiability in β of their limit functions, that are not needed here. Similarly, their conditions almost require $\max_{s \leq L} \mu_n(s) \rightarrow_p 0$ where we come away with pointwise convergence. (iii) is interesting to see that the key requirement (6.11) serves two different purposes: forcing an analytical remainder term towards zero as well as securing uniform negligibility of individual terms, i.e. limiting normality. (iv) The methods used here can be applied to solve the large-sample behaviour problem also outside model conditions, say when the true hazard rate is $\lambda(s) r(z_{i,1}(s), \dots, z_{i,p}(s))$ for individual i . See Hjort (1992) for results. There are also various alternative estimation techniques that can be employed in the Cox model, see for examples Hjort (1991) for local likelihood smoothing and Hjort (1992) for weighted log partial likelihood estimation. Again techniques from the present paper can be applied. (v) Finally Jeffreys type arguments can be given in favour of using the vague prior $\pi(\beta) = 1$, see Hjort (1986), where it is also shown that the (improper) pseudo-Bayes estimator $\beta_n^* = \int \beta \exp(G_n(\beta)) d\beta / \int \exp(G_n(\beta)) d\beta$ is asymptotically equivalent to the Cox estimator $\hat{\beta}_n$. The arguments of 4B can be used to provide a quicker and simpler proof of this.

7. Further applications.

7A. *Exponential hazard rate regression.* The traditional Cox model (6.1) is semiparametric, since the basis hazard rate $\lambda(\cdot)$ there is left unspecified. The parametric version

$$\lambda_i(s) = \lambda_0(s) \exp(\beta' z_i(s)) = \lambda_0(s) \exp(\beta_1 z_{i,1}(s) + \dots + \beta_p z_{i,p}(s)), \quad (7.1)$$

where $\lambda_0(\cdot)$ is fully specified (equal to 1, for example), is also important in survival data analysis. Let us briefly show that the arguments above efficiently lead to a precise theorem about the maximum likelihood estimator in this model as well.

With notation and assumptions otherwise being as in section 6 the log-likelihood can be written

$$\log L_n(\beta) = \sum_{i \leq n} \int_0^L \{ \beta' z_i(s) dN_i(s) - Y_i(s) \exp(\beta' z_i(s)) ds \},$$

see for example Andersen et al. (1992, chapter VI), and let $\hat{\beta}_n$ be the ML estimator maximising this expression. Assume that data follow (7.1) for a certain β_0 . Using martingales $dM_i(s) = dN_i(s) - Y_i(s) \exp(\beta_0' z_i(s)) d\Lambda_0(s)$, writing $d\Lambda_0(s) = \lambda_0(s) ds$, we find

$$\begin{aligned} G_n(x) &= \log L_n(\beta_0 + x) - \log L_n(\beta_0) \\ &= \sum_{i \leq n} \int_0^L \left[x' z_i(s) \{ dM_i(s) + Y_i(s) \exp(\beta_0' z_i(s)) d\Lambda_0(s) \} \right. \\ &\quad \left. - Y_i(s) \exp(\beta_0' z_i(s)) \{ \exp(x' z_i(s)) - 1 \} ds \right] \\ &= \left(\sum_{i \leq n} \int_0^L z_i(s) dM_i(s) \right)' x - \sum_{i \leq n} \int_0^L Y_i(s) \exp(\beta_0' z_i(s)) \{ \exp(x' z_i(s)) - 1 - x' z_i(s) \} d\Lambda_0(s). \end{aligned}$$

Now use $|\exp(u) - 1 - u - \frac{1}{2}u^2| \leq \frac{1}{6}|u|^3 \exp(|u|)$, and introduce

$$J_n = \sum_{i \leq n} \int_0^L Y_i(s) \exp(\beta_0' z_i(s)) z_i(s) z_i(s)' d\Lambda_0(s) \text{ and } U_n = J_n^{-1/2} \sum_{i \leq n} \int_0^L z_i(s) dM_i(s) \quad (7.2)$$

to find $G_n(J_n^{-1/2}x) = U_n'x - \frac{1}{2}|x|^2 - r_n(x)$, with a remainder bound

$$|r_n(x)| \leq \frac{1}{6} \sum_{i \leq n} \int_0^L Y_i(s) \exp(\beta_0' z_i(s) + |x' J_n^{-1/2} z_i(s)|) \frac{1}{6} |x' J_n^{-1/2} z_i(s)|^3 ds. \quad (7.3)$$

To formulate a theorem with quite weak conditions, let $J_n(s) = \sum_{i \leq n} Y_i(s) \exp(\beta_0' z_i(s)) z_i(s) z_i(s)'$, so that the ‘observed information matrix’ is $J_n = \int_0^L J_n(s) d\Lambda_0(s)$.

THEOREM 7.1. *Let $\lambda_0(s)$ be positive and continuous on $[0, L]$. Suppose there is a c_n sequence converging to infinity such that $J_n(s)/c_n$ for almost all s goes in probability to some $J(s)$, and that $J_n/c_n \rightarrow_p J = \int_0^L J(s) d\Lambda_0(s)$, where this limit matrix is positive definite. Assume furthermore that for almost all s ,*

$$N_n(s, \delta) = \sum_{i \leq n} z_i(s)' J_n^{-1} z_i(s) Y_i(s) \exp(\beta_0' z_i(s)) I\{|J_n^{-1/2} z_i(s)| \geq \delta\} \rightarrow_p 0 \quad (7.4)$$

for each $\delta > 0$, and that $\mu_n(s) = \max_{i \leq n} |J_n^{-1/2} z_i(s)|$ is stochastically bounded, uniformly in s . Then $J_n^{1/2}(\hat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_p\{0, I_p\}$.

Some brief remarks are in order before turning to the proof. (i) $z_i(s) J_n^{-1} z_i(s)$ can be replaced by $z_i(s)' J^{-1} z_i(s)/c_n$ here, and $|J_n^{-1/2} z_i(s)|$ with $|J^{-1/2} z_i(s)|/c_n^{1/2}$. (ii) In many practical situations the c_n will be equal to n . (iii) The elements of J_n may in some cases conceivably go to infinity with different rates, and then the ‘asymptotic stability’ requirement should be the existence of matrices C_n going to infinity such that $C_n^{-1} J_n(s) \rightarrow J(s)$ etcetera. The theorem still holds. (iv) In many cases one would have $\mu_n(s) \rightarrow_p 0$ for almost all s , and this implies condition (7.4), since in fact $D_n(s, \delta) \leq p I\{\mu_n(s) \geq \delta\}$. (v) If the $z_i(s)$ covariate processes are uniformly bounded, then (iv) applies and hence the conclusion. (vi) Our conditions are much weaker than those used elsewhere to secure large sample normality, see for example Borgan (1984, section 6). (vii) Finally we note that the proof below becomes easier under circumstances (iv) or (v).

PROOF: The log-likelihood is concave by Lemma A.2 and hence so is the $G_n(J_n^{-1/2}x)$ function. We are to prove (i) $r_n(x) \rightarrow 0$ in probability for each x , and (ii) that $U_n \rightarrow \mathcal{N}_p\{0, I_p\}$ in distribution.

To prove (i) let $r_n(x, s)$ be the integrand in the bound occurring in (7.3), so that $|r_n(x)| \leq \frac{1}{6} \int_0^L r_n(x, s) d\Lambda_0(s)$. It will suffice to show that $r_n(x, s) \rightarrow 0$ in probability for almost all s and to bound it properly. Splitting into $|J_n^{-1/2} z_i(s)| < \delta$ terms and $|J_n^{-1/2} z_i(s)| \geq \delta$ terms we find $r_n(x, s) \leq |x|^3 \delta \exp(|x|\delta) + |x|^3 \mu_n(s) \exp(|x|\mu_n(s)) N_n(s, \delta)$, after which the claim follows by our precautions and by the dominated convergence lemma of the appendix.

Next (ii) can be replaced by $U_n^* \rightarrow_d \mathcal{N}_p\{0, I_p\}$, where $U_n^* = \sum_{i \leq n} \int_0^L c_n^{-1/2} J^{-1/2} z_i(s) dM_i(s)$, and we show this employing the Rebolledo theorem version given in Andersen and Gill (1982, appendix I). Its variance process converges properly,

$$\begin{aligned} \langle U_n^*, U_n^* \rangle(L) &= \sum_{i \leq n} \int_0^L c_n^{-1} J^{-1/2} z_i(s) z_i(s)' J^{-1/2} Y_i(s) \exp(\beta_0' z_i(s)) d\Lambda_0(s) \\ &= J^{-1/2} (J_n/c_n) J^{-1/2} \rightarrow_p I_p, \end{aligned}$$

and the Lindebergian condition is also satisfied:

$$\sum_{i \leq n} \int_0^L c_n^{-1} |J^{-1/2} z_i(s)|^2 I\{c_n^{-1/2} |J_n^{-1/2} z_i(s)| \geq \delta\} Y_i(s) \exp(\beta_0' z_i(s)) d\Lambda_0(s) \rightarrow_p 0.$$

This holds since the integrand is asymptotically the same as $N_n(s, \delta)$ of (7.4), and is bounded by the constant p , so that the dominated convergence lemma applies. \square

7B. *Poisson regression.* Suppose Y_1, \dots, Y_n are independent counts with

$$Y_i \sim \text{Poisson}(\text{mean}_i), \quad \text{with } \text{mean}_i = \exp(\beta' z_i), \quad (7.5)$$

depending on a certain p -dimensional covariate vector z_i , for a certain true parameter value β_0 . It is convenient now to write $\exp(u) = 1 + u + \frac{1}{2}u^2 + \frac{1}{6}\rho(u)$, where a bound for the remainder function is $|\rho(u)| \leq |u|^3 \exp(|u|)$. The log-likelihood $\log L_n(\beta) = \sum_{i \leq n} \{Y_i \beta' z_i - \exp(\beta' z_i)\}$ is concave, and after development very similar to that of section 7A one finds

$$\log L_n(\beta_0 + x) - \log L_n(\beta_0) = \left(\sum_{i \leq n} (Y_i - \mu_i) z_i \right)' x - \frac{1}{2} x' J_n x - \frac{1}{6} v_n(x),$$

in which $J_n = \sum_{i \leq n} \mu_i z_i z_i'$ and $v_n(x) = \sum_{i \leq n} \mu_i \rho(x' z_i)$. In these expressions $\mu_i = \exp(\beta_0' z_i)$ is the true mean for Y_i under the model.

Before passing to a theorem we solve a relevant exercise in asymptotics of linear combinations of independent Poisson variables. If $Y_{n,i}$ is Poisson with mean $\mu_{n,i}$, then $\sum_{i \leq n} (Y_{n,i} - \mu_{n,i}) x_{n,i}$, normed such that its variance $\sum_{i \leq n} \mu_{n,i} x_{n,i}^2 = 1$, goes to a standard normal if and only if $\sum_{i \leq n} \mu_{n,i} \rho(t x_{n,i}) \rightarrow 0$ for each t , which is equivalent to $\sum_{i \leq n} \mu_{n,i} \rho(|x_{n,i}|) \rightarrow 0$. This is seen after considering moment or cumulant generating functions.

THEOREM 7.2. *Let $\hat{\beta}_n$ be the ML estimator based on the first n Poisson counts, assumed to follow (7.5) for a certain β_0 , with means $\mu_i = \exp(\beta_0' z_i)$. Then $J_n^{1/2}(\hat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}_p\{0, I_p\}$ if and only if $\sum_{i \leq n} \mu_i \rho(|J_n^{-1/2} z_i|) \rightarrow 0$. A simple sufficient condition for this to hold is that $\lambda_n = \max_{i \leq n} |J_n^{-1/2} z_i|$ is bounded and that $\sum_{i \leq n} \mu_i |J_n^{-1/2} z_i|^3 \rightarrow 0$; or, equivalently, that λ_n is bounded and that*

$$N_n(\delta) = \sum_{i \leq n} \mu_i z_i' J_n^{-1} z_i I\{|J_n^{-1/2} z_i| \geq \delta\} \rightarrow 0 \quad \text{for each } \delta.$$

PROOF: The function $\log L_n(\beta_0 + J_n^{-1/2} x) - \log L_n(\beta_0)$ is concave in x and can be written $U_n' x - \frac{1}{2}|x|^2 - r_n(x)$, where $U_n = J_n^{-1/2} \sum_{i \leq n} (Y_i - \mu_i) z_i$ and where $r_n(x) = \sum_{i \leq n} \mu_i \rho(x' J_n^{-1/2} z_i)$. This is quite similar to the situation in 7A, and the maximiser $J_n^{1/2}(\hat{\beta}_n - \beta_0)$ goes to a standard p -dimensional normal if and only if (i) $r_n(x) \rightarrow 0$ and (ii) $U_n \rightarrow_d \mathcal{N}_p\{0, I_p\}$. But using the result above in tandem with the Cramér–Wold theorem one sees that $\sum_{i \leq n} \mu_i \rho(|J_n^{-1/2} z_i|) \rightarrow 0$ is necessary and sufficient for (ii), and indeed also necessary and sufficient for (i). The other statements of the theorem follow from $|\rho(u)| \leq |u|^3 \exp(|u|)$. \square

We note that $\lambda_n \rightarrow 0$ is clearly sufficient for the result to hold.

7C. *Generalised linear models.* Consider a situation with independent Y_i 's from densities of the form $f(y_i | \theta_i) = \exp\{(y_i \theta_i - b(\theta_i))/a(\phi) + c(y_i, \phi)\}$, and where θ_i is parametrised as a linear $x_i' \beta$. This is a generalised linear model with canonical link, see McCullagh and Nelder (1989). The likelihood in β is log-concave, and theorems about the large-sample behaviour of the ML estimator, under very weak regularity conditions, can be written down and proved by the methods exemplified in sections 5 and 7A.

too much?

7D. Pseudo-likelihood estimation in Markov chains. Suppose X_0, X_1, \dots forms a Markov chain on the state space $\{1, \dots, k\}$. Instead of focusing on transition probabilities, consider direct modelling of X_i given its neighbours, say $X_{\partial i} = x_{\partial i}$. A very flexible and convenient class of models is described by

$$f_{\beta}(x_i|x_{\partial i}) = \text{const.} \exp\{\alpha_i(x_i) + \beta' H_i(x_i, x_{\partial i})\} = \frac{\exp\{\alpha_i(x_i) + \beta' H_i(x_i, x_{\partial i})\}}{\sum_{j=1}^k \exp\{\alpha_i(j) + \beta' H_i(j, x_{\partial i})\}}, \quad (7.6)$$

where $\alpha_i(1), \dots, \alpha_i(k)$ are specified or unknown parameters, and $\beta' H_i(x_i, x_{\partial i}) = \sum_{u=1}^p \beta_u H_{i,u}(x_i, x_{\partial i})$ for certain component functions $H_{i,u}$ that depend both on the x_i at time position i and of its neighbouring values $x_{\partial i}$. For a second order Markov chain, for example, one would typically have H_i equal to a common H function for $2 \leq i \leq n-2$ and some special functions at the borders. Maximum pseudo-likelihood estimation maximises $\text{PL}_n(\beta) = \prod_{i=0}^n f_{\beta}(x_i|x_{\partial i})$ w.r.t. the parameters. See Hjort and Omre (1993, section 3.2), for example, for comments on this model building machinery in dimensions 1 and 2, and for some comments on the difference between maximum PL and maximum likelihood.

We may incorporate the $\alpha_i(x)$'s in the vector $H_i(x, x_{\partial i})$ of $H_{i,u}$ -functions, for notational convenience. From Lemma A2 $\log \text{PL}_n(\beta)$ is concave in β . Consider $h_i(x_{\partial i}) = \sum_{j=1}^k H_i(j, x_{\partial i}) f_{\beta_0}(j|x_{\partial i})$ and

$$V_i(x_{\partial i}) = \sum_{j=1}^k (H_i(j, x_{\partial i}) - h_i(x_{\partial i})) (H_i(j, x_{\partial i}) - h_i(x_{\partial i}))' f_{\beta_0}(j|x_{\partial i}),$$

which can be interpreted as respectively $E\{H_i(X_i, x_{\partial i})|x_{\partial i}\}$ and $\text{VAR}\{H_i(X_i, x_{\partial i})|x_{\partial i}\}$. After some work exploiting Lemma A2 one finds that

$$\log \text{PL}_n(\beta_0 + s/\sqrt{n}) - \log \text{PL}_n(\beta_0) = U_n' s - \frac{1}{2} s' (J_n/n) s - r_n(s), \quad (7.7)$$

where

$$U_n = n^{-1/2} \sum_{i=0}^n \{H_i(X_i, X_{\partial i}) - h_i(X_{\partial i})\} \quad \text{and} \quad J_n = \sum_{i=0}^n V_i(X_{\partial i}),$$

and where in fact $r_n(s) = O(n^{-1/2})$. The usual arguments now give $\sqrt{n}(\hat{\beta}_n - \beta_0) \rightarrow_d \mathcal{N}\{0, J^{-1}\}$ under mild assumptions, provided the assumed model (7.6) is correct. Here J turns out to be both the limit of J_n/n as well as the covariance matrix in the limiting distribution for U_n . There is also an appropriate sandwich generalisation with covariance matrix of type $J^{-1} K J^{-1}$ outside model conditions. Doing the details here properly calls for a central limit theorem and a weak law of large numbers for Markov chains, and such can be found in Billingsley (1961), for example.

These Markov random field models are more important in the 2- and 3-dimensional cases, where one enters the world of statistical image analysis. The method above can be used to prove consistency of the maximum PL estimator.

Appendix. Here we give three lemmas that were used at various stages above. They should also have some independent interest.

A1. Necessary and sufficient conditions for asymptotic normality of linear combinations of binomials. The following result with further consequences was used in section 5.

LEMMA A1. Consider independent Bernoulli variables $Y_{n,i} \sim \text{Bin}\{1, q_{n,i}\}$, and real numbers $z_{n,i}$ standardised to have $\sum_{i \leq n} z_{n,i}^2 q_{n,i} (1 - q_{n,i}) = 1$. Then $\sum_{i \leq n} z_{n,i} (Y_{n,i} - q_{n,i}) \rightarrow_d \mathcal{N}\{0, 1\}$ if and only if

$$N_n(\delta) = \sum_{i \leq n} z_{n,i}^2 q_{n,i} (1 - q_{n,i}) I\{|z_{n,i}| \geq \delta\} \rightarrow 0 \quad \text{for each positive } \delta. \quad (\text{A.1})$$

PROOF: The Lindeberg condition is that

$$\begin{aligned} L_n(\delta) &= \sum_{i \leq n} E z_{n,i}^2 (Y_{n,i} - q_{n,i})^2 I\{|z_{n,i}(Y_{n,i} - q_{n,i})| \geq \delta\} \\ &= \sum_{i \leq n} z_{n,i}^2 q_{n,i}(1 - q_{n,i}) [q_{n,i} I\{|q_{n,i} z_{n,i}| \geq \delta\} + (1 - q_{n,i}) I\{|(1 - q_{n,i}) z_{n,i}| \geq \delta\}] \end{aligned}$$

should tend to zero for each positive δ . It is not difficult to establish $\frac{1}{2}N_n(2\delta) \leq L_n(\delta) \leq N_n(\delta)$, so (A.1) is in fact equivalent to the Lindeberg requirement. In particular (A.1) implies a $\mathcal{N}\{0, 1\}$ limit.

Necessity is harder. Assume a $\mathcal{N}\{0, 1\}$ limit in distribution. We first symmetrise in the following fashion: Let $\tilde{Y}_{n,i} = Y_{n,i} - Y'_{n,i}$ where $Y'_{n,1}, Y'_{n,2}, \dots$ are independent copies of $Y_{n,1}, Y_{n,2}, \dots$, and let

$$Z_n = \sum_{i \leq n} z_{n,i}(Y_{n,i} - q_{n,i}), \quad Z'_n = \sum_{i \leq n} z_{n,i}(Y'_{n,i} - q_{n,i}), \quad \text{and} \quad \tilde{Z}_n = Z_n - Z'_n.$$

By assumption $\tilde{Z}_n \rightarrow_d \mathcal{N}\{0, 2\}$. We first show that

$$m_n = \max_{i \leq n} \min\{|z_{n,i}|, q_{n,i}, 1 - q_{n,i}\} \rightarrow 0.$$

Otherwise there would be some $\varepsilon > 0$ such that say $|z_{n,1}| \geq \varepsilon$ and $\varepsilon \leq q_{n,1} \leq 1 - \varepsilon$. Break \tilde{Z}_n into a sum of $V_n = z_{n,1}\tilde{Y}_{n,1}$ and W_n , two independent and symmetric variables. Uniform tightness of \tilde{Z}_n and symmetry imply uniform tightness of both V_n and W_n . Along some subsequence we would have $V_n \rightarrow_d V$ and $W_n \rightarrow_d W$, independent with $V + W$ distributed as $\mathcal{N}\{0, 2\}$. By Cramér's theorem about convolution factors of $\mathcal{N}\{0, 2\}$ we would have V normal. But V is not degenerate, and cannot be normal after all, since V_n takes only three values. This proves $m_n \rightarrow 0$.

But this implies the usual infinitesimal array property

$$\max_{i \leq n} \Pr\{|z_{n,i}\tilde{Y}_{n,i}| \geq \delta\} \rightarrow 0 \quad \text{for each } \delta.$$

For if $|z_{n,i}| < \delta$ then the probability is zero, and if $|z_{n,i}| \geq \delta$ then the probability is $2q_{n,i}(1 - q_{n,i}) \leq 2m_n$ when n is large enough for $m_n < \delta$ to hold. Next look at page 92 of Petrov (1975). From limiting normality follows

$$\sum_{i \leq n} \text{Var} [z_{n,i}\tilde{Y}_{n,i} I\{|z_{n,i}\tilde{Y}_{n,i}| < \delta\}] \rightarrow 2.$$

If $|z_{n,i}| \geq \delta$ the indicator here picks out $\tilde{Y}_{n,i} = 0$, and there is no contribution to the sum, whereas if $|z_{n,i}| < \delta$ the summand is $2z_{n,i}^2 q_{n,i}(1 - q_{n,i})$. Hence $\sum_{i \leq n} z_{n,i}^2 q_{n,i}(1 - q_{n,i}) I\{|z_{n,i}| < \delta\} \rightarrow 1$, and $N_n(\delta) \rightarrow 0$ follows from the assumed $\sum_{i \leq n} z_{n,i}^2 q_{n,i}(1 - q_{n,i}) = 1$. \square

The surprising thing here is that we do not need to explicitly assume $\max_{i \leq n} E\{z_{n,i}(Y_{n,i} - q_{n,i})\}^2 \rightarrow 0$, as with Feller's partial converse to the Lindeberg theorem; it follows from asymptotic normality and the special properties of the $Y_{n,i}$ sequence.

Lemma A1 can next be used to address the vector case, via the Cramér–Wold theorem. We phrase the result as follows, to suit the development of section 5. If x_1, x_2, \dots is a sequence of p -vectors, and Y_1, Y_2, \dots are Bernoulli with q_1, q_2, \dots , then

$$J_n^{-1/2} \sum_{i \leq n} (Y_i - q_i)x_i \rightarrow_d \mathcal{N}_p\{0, I_p\}, \quad \text{where } J_n = \sum_{i \leq n} q_i(1 - q_i)x_i x_i', \quad (\text{A.2})$$

if and only if

$$N_n(\delta) = \sum_{i \leq n} x'_i J_n^{-1} x_i q_i (1 - q_i) I\{|J_n^{-1/2} x_i| \geq \delta\} \rightarrow 0 \quad \text{for each } \delta > 0. \quad (\text{A.3})$$

This is proved by noting first that (A.2) is equivalent to

$$N_n^0(s, \delta) = \sum_{i \leq n} s' x'_i J_n^{-1} x_i s q_i (1 - q_i) I\{|s' J_n^{-1/2} x_i| \geq \delta\} \rightarrow 0$$

for all s with length 1 and all positive δ . But $N_n^0(s, \delta) \leq N_n(\delta) \leq p^2 \max_{j \leq p} N_n^0(e_j, \delta/\sqrt{p})$, where e_j is the j th unit vector.

A simple *sufficient* condition for (A.1) to hold is that $\lambda_n^0 = \max_{i \leq n} |z_{n,i}| \rightarrow 0$, since the left hand side of (A.1) is bounded by λ_n^0/δ . Similarly condition (A.3) is implied by the simpler condition $\lambda_n = \max_{i \leq n} |J_n^{-1/2} x_i| \rightarrow 0$, since $N_n(\delta) \leq \lambda_n p/\delta$.

A2. Expansion lemma. The following result was used in section 6 and in 7A.

LEMMA A2. (i) Suppose $K(t) = \log R(t)$, where $R(t) = \sum_{i \leq n} w_i \exp(a_i t)$ for certain nonnegative weights w_i , not all equal to zero, and arbitrary constants a_i . Let $v_i(t) = w_i \exp(a_i t)/R(t)$ be the tilted and normalised weights, summing to one. Then $K(t)$ is convex with derivatives

$$\begin{aligned} K'(t) &= \sum_{i \leq n} v_i(t) a_i = \bar{a}(t), \\ K''(t) &= \sum_{i \leq n} v_i(t) \{a_i - \bar{a}(t)\}^2, \\ K'''(t) &= \sum_{i \leq n} v_i(t) \{a_i - \bar{a}(t)\}^3. \end{aligned}$$

(ii) *The expansion*

$$\log \left\{ \sum_{i \leq n} w_i e^{a_i t} \right\} - \log \left\{ \sum_{i \leq n} w_i \right\} = \bar{a}(0)t + \frac{1}{2} \sum_{i \leq n} v_i(0) \{a_i - \bar{a}(0)\}^2 t^2 + v(t)$$

holds, featuring untilted weights $v_i(0) = w_i/\sum_{i \leq n} w_i$, with the following valid bounds on the remainder:

$$|v(t)| \leq \frac{4}{3} \mu_n^3 |t|^3, \quad |v(t)| \leq \frac{2}{3} g(\mu_n |t|) \sum_{i \leq n} v_i(0) \{a_i - \bar{a}(0)\}^2 |t|^2. \quad (\text{A.4})$$

Here $\mu_n = \max_{i \leq n} |a_i - \bar{a}(0)|$ and g is the function $g(u) = u \exp(2u + 4u^2)$.

PROOF: The formulae for the derivatives are proved by direct differentiation and inspection, and convexity follows of course from the nonnegative second derivative. To prove (ii), consider the exact third order Taylor expansion $K(t) - K(0) = K'(0)t + \frac{1}{2} K''(0)t^2 + \frac{1}{6} K'''(s)t^3$ for some suitable s between 0 and t . The problem is to bound the remainder term in terms of μ_n .

The first bound is easy. It follows upon observing that $|\bar{a}(s) - \bar{a}(0)| = |\sum_{i \leq n} v_i(s) \{a_i - \bar{a}(0)\}| \leq \mu_n$ and its triangle inequality consequence $|a_i - \bar{a}(s)| \leq 2\mu_n$, since this yields $|K'''(s)| \leq (2\mu_n)^3$. While this bound often suffices we shall have occasion to need the sharper second bound too. The point is to exploit the fact that s is bounded by $|t|$ when bounding $|K'''(s)|$. Start out writing $v_i(s) = v_i(0)(1 + \varepsilon_i)$, where some analysis shows that $\exp(-2\mu_n |t|) \leq 1 + \varepsilon_i \leq \exp(2\mu_n |t|)$. Then

$$\begin{aligned} |\bar{a}(s) - \bar{a}(0)| &= \left| \sum_{i \leq n} v_i(0)(1 + \varepsilon_i) \{a_i - \bar{a}(0)\} \right| \\ &= \left| \sum_{i \leq n} v_i(0) \varepsilon_i \{a_i - \bar{a}(0)\} \right| \leq K''(0)^{1/2} \left\{ \sum_{i \leq n} v_i(0) \varepsilon_i^2 \right\}^{1/2}. \end{aligned}$$

A further bound on the right hand side is $K''(0)^{1/2}\delta_n = K''(0)^{1/2} \max_{i \leq n} |\varepsilon_i|$. This gives

$$\begin{aligned} |K'''(s)| &\leq 2\mu_n \sum_{i \leq n} v_i(s) |a_i - \bar{a}(s)|^2 \\ &\leq 4\mu_n \sum_{i \leq n} v_i(0) (1 + \varepsilon_i) \{|a_i - \bar{a}(0)|^2 + |\bar{a}(0) - \bar{a}(s)|^2\} \\ &\leq 4\mu_n (1 + \delta_n) (1 + \delta_n^2) K''(0). \end{aligned}$$

But some checking reveals $1 + \delta_n \leq \exp(2\mu_n|t|)$ and $1 + \delta_n^2 \leq \exp(4\mu_n^2|t|^2)$. This shows $|K'''(s)| \leq 4|t|^{-1}g(\mu_n|t|)K''(0)$ with the g -function given above. \square

A3. Dominated convergence theorem for convergence in probability. This result was used several times in section 6, and a close relative was used in 4B.

LEMMA A3. *Let $0 \leq X_n(s, \omega) \leq Y_n(s, \omega)$ be jointly (s, ω) -measurable random functions on the interval $[0, L]$. Suppose λ is a measure such that $Y_n(s) \rightarrow_p Y(s)$ and $X_n(s) \rightarrow_p X(s)$ for λ almost all s and that $\int Y_n(s) d\lambda(s) \rightarrow_p \int Y(s) d\lambda(s)$, a limit finite almost everywhere. Then $\int X_n(s) d\lambda(s) \rightarrow_p \int X(s) d\lambda(s)$ too.*

PROOF: It is enough to check almost sure convergence for a subsubsequence of each subsequence. By convergence in probability (for ω , with s fixed) and then dominated convergence, we have

$$\pi_n(\varepsilon) := (\mathbb{P} \otimes \lambda) \{(\omega, s) : |X_n(\omega, s) - X(\omega, s)| > \varepsilon\} \rightarrow 0.$$

A similar result holds for $\{Y_n\}$. Replace ε by a sequence $\{\varepsilon_n\}$ decreasing to zero, then extract a subsubsequence along which the sequence of integrals is convergent. For some set N with $(\mathbb{P} \otimes \lambda)N = 0$ we get convergence for all $(\omega, s) \in N^c$ of both the X_n and the Y_n subsubsequences. For almost all ω , therefore, $\lambda\{s : (\omega, s) \in N\} = 0$. Finally argue using the Fatou Lemma for $Y_n \pm X_n$ along the subsubsequences to get the result. \square

This is nice in that it circumvents the need to establish uniformity of the convergence in probability; this is typically more difficult to ascertain than pointwise convergence in probability. The lemma was used several times in sections 6 and 7A, partly in the form of the following useful corollary: If in particular $Z_n(s) \rightarrow_p 0$ for almost all s , then $\int_0^L Y_n(s) I\{|Z_n(s)| \geq \delta\} ds \rightarrow_p 0$. It can also be used to simplify the Lindeberg type condition in the form of Rebolledo's martingale convergence theorem given in Andersen and Gill (1982, appendix).

Acknowledgements. This paper resulted from a collaboration made possible by the support of the Mathematical Sciences Research Institute at Berkeley during the Autumn of 1991, under National Science Foundation grant 8505550. We are grateful to the MSRI for the opportunity to learn from each other. N.L.H. has also been partly supported by grants from the Royal Norwegian Research Council and D.P. partly supported by National Science Foundation Grant DMS-9102286.

References

- Andersen, P.K., Borgan, Ø., Gill, R.D., and Keiding, N. (1992). *Statistical Models Based on Counting Processes*. Springer-Verlag, New York.
- Andersen, P.K. and Gill, R.D. (1982). Cox's regression model for counting processes: a large sample study. *Ann. Statist.* **10**, 1100–1120.
- Bassett, G.W. and Koenker, R.W. (1982). An empirical quantile method for linear models with iid errors. *J. Amer. Statist. Assoc.* **77**, 407–415.

- Bickel, P.J., Klaassen, C.A., Ritov, Y., and Wellner, J.A. (1992). *Efficient and Adaptive Inference in Semiparametric Models*. To exist.
- Billingsley, P. (1961). *Statistical Inference for Markov Processes*. University of Chicago Press, Chicago.
- Borgan, Ø. (1984). Maximum likelihood estimation in parametric counting process models, with applications to censored failure time data. *Scand. J. Statist.* **11**, 1–16. [Corrigendum, *ibid.* p. 275.]
- Haberman, S.J. (1989). Concavity and estimation. *Ann. Statist.* **17**, 1631–1661.
- Hjort, N.L. (1986). Bayes estimators and asymptotic efficiency in parametric counting process models. *Scand. J. Statist.* **13**, 63–85.
- Hjort, N.L. (1988a). Logistic regression when the model is wrong. Appendix to *Statistical models for the probability of finding oil or gas*, with V. Berteig. Norwegian Computing Centre Report, Oslo.
- Hjort, N.L. (1988b). Lecture notes and seventy-five exercises on resampling and bootstrapping. Statistical Research Report, University of Oslo.
- Hjort, N.L. (1991). Semiparametric estimation of parametric hazard rates. In *Survival Analysis: State of the Art*, Kluwer, Dordrecht, pp. 211–236. Proceedings of the *NATO Advanced Study Workshop on Survival Analysis and Related Topics*, Columbus, Ohio, eds. P.S. Goel and J.P. Klein.
- Hjort, N.L. (1992). On inference in parametric survival data models. *Intern. Statist. Review* **60**, 355–387.
- Hjort, N.L. and Omre, H. (1993). Topics in spatial statistics (with discussion). *Scand. J. Statist.*, to appear.
- Huber, P. (1967). The behaviour of maximum likelihood estimates under non-standard conditions. *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability* **1**, 221–233. University of California Press.
- Jurečková, J. (1977). Asymptotic relations of M-estimates and R-estimates in linear regression model. *Ann. Statist.* **5**, 464–472.
- Jurečková, J. (1992). Estimation in a linear model based on regression rank scores. *J. Nonparametric Statistics* **1**, 197–203.
- Lehmann, E.L. (1983). *Theory of Point Estimation*. Wiley, New York.
- McCullagh, P. and Nelder, J.A. (1989). *Generalized Linear Models* (2nd edition). Chapman & Hall, London.
- Niemiro, W. (1992). Asymptotics for M-estimators defined by convex minimization. *Ann. Statist.* **20**, 1514–1533.
- Petrov, V.V. (1975). *Sums of Independent Random Variables*. Springer-Verlag, New York.
- Pollard, D. (1984). *Convergence of Stochastic Processes*. Springer-Verlag, New York.
- Pollard, D. (1985). New ways to prove central limit theorems. *Econometric Theory* **1**, 295–314.
- Pollard, D. (1990). *Empirical Processes: Theory and Applications*. NSF-CBMS Regional Conference Series in Probability and Statistics, Vol. 2. IMS, Hayward, California.
- Pollard, D. (1991). Asymptotics for least absolute deviation regression estimators. *Econometric Theory* **7**, 186–199.
- Rockafellar, R.T. (1970). *Convex Analysis*. Princeton University Press, Princeton, New Jersey.